

Original Article

A Systematic Review of Alveolar Defect Healing Following Tooth Extraction through Primary and Secondary Intention

Razukevicius Dainius¹, Janovskiene Audra^{1*}, Micka Kornelijus², Petronis Zygimantas¹, Chomicius Deividas², Daugela Povilas¹

¹ Department of Maxillofacial Surgery, Lithuanian University of Health Sciences, Kaunas Lithuania.

² Department of Odontology, Lithuanian University of Health Sciences, Kaunas Lithuania.

*E-mail 🖂 a.janovskiene@gmail.com

Received: 19 February 2025; Revised: 20 May 2025; Accepted: 21 May 2025

ABSTRACT

Tooth extraction leads to changes in the dimensions of the alveolar ridge, prompting efforts to reduce complications and bone loss. Various strategies have been investigated to preserve bone width, height, and keratinized tissue after extractions. This systematic review analyzes the effect of secondary and primary intention healing on the dimensional changes and histomorphometric properties of the alveolar ridge. In adherence to PRISMA guidelines, a literature search was performed on the Cochrane Library, PubMed, and ScienceDirect databases. This review included randomized controlled trials, retrospective studies, observational studies, split-mouth randomized trials, single-blinded, and controlled clinical trials. Articles published between March 1, 2013, and March 1, 2023, that compared bone changes or histomorphometric data after tooth extraction were included. The risk of bias was assessed using the "Cochrane Collaboration's risk-of-bias (RoB 2) tool." A total of 5 studies, involving 92 patients and 128 extracted teeth, were reviewed. The findings showed no significant difference in the resorption of alveolar height and width between the groups treated with primary or secondary intention. Similarly, there were no significant differences in histomorphometric changes between the two healing methods. Therefore, the healing method choice did not significantly affect the dimensional alterations or new bone formation in post-extraction defects.

Keywords: Socket, Primary intention, Open flap, Secondary intention, Closed flap.

How to Cite This Article: Dainius R, Audra J, Kornelijus M, Zygimantas P, Deividas C, Povilas D. A Systematic Review of Alveolar Defect Healing Following Tooth Extraction through Primary and Secondary Intention. J Curr Res Oral Surg. 2025;5:44-51. https://doi.org/10.51847/acsCjm1kLP

Introduction

After tooth extraction, the alveolar ridge undergoes changes that present challenges for dental specialists in daily clinical practice. The ridge's dimensions, including its shape and size, are influenced by the tooth's structure and the vertical and horizontal alterations that follow tooth loss [1]. Around 60% of the ridge volume decreases within the first 2–3 years post-extraction, with the resorption continuing at a rate of 0.25–0.5% annually throughout life [2]. Most of this

bone loss occurs within the first few months after extraction [3], and horizontal bone loss tends to be more significant than vertical loss [4]. Research shows that the outer (buccal) part of the alveolar ridge experiences more resorption than the inner (lingual) side [5]. This is due to the thinner buccal wall, which makes it more susceptible to dehiscence [6]. When there are fewer bony walls, there is less opportunity to retain a blood clot in the socket [7]. Several factors, including the thickness of the alveolar wall, tooth position, surgical trauma, flap elevation, and the size of

the original defect, determine the extent of bone resorption following extraction [8]. Studies indicate that about 30% of the alveolar ridge undergoes resorption after tooth extraction, with approximately two-thirds of both soft and hard tissues being affected within the first three months [9]. To minimize bone loss during the healing process, researchers are focused on methods and materials that can help preserve or restore bone width, height, and keratinized tissues.

After tooth extraction, various materials, including autogenous, allogenic, xenogenic, and alloplastic bone granules, are employed to maintain the socket's dimensions and encourage the regeneration of viable bone tissue. The selection of these materials may be influenced by the clinician's preference, as well as financial or cultural factors [7]. To stabilize the bone grafts, different membranes are used, ranging from resorbable options like collagen films to nonresorbable membranes, such as PTFE or titanium, which require an additional surgical procedure for removal. Several techniques are available for closing the post-extraction wound to shield the socket from external threats. In primary intention healing, the wound is sealed using a mucoperiosteal flap, while in secondary intention healing, the flap is not completely sutured, allowing for drainage or sometimes left unsutured [10, 11]. The majority of the scientific literature suggests that to prevent infection, most membranes should be entirely covered with a mobilized mucoperiosteal flap for primary intention healing. Nevertheless, it is also possible to leave certain membranes, like d-PTFE or collagen, without suturing for secondary intention healing [10].

This study aims to evaluate the effects of primary and secondary intention healing on the dimensional changes of the alveolar ridge and histomorphometric bone tissue markers following tooth extraction.

Materials and Methods

In conducting this systematic review, the study adhered to the PRISMA guidelines, which guided the planning, objectives, article selection, and data analysis process [12]. The PICO framework was used to structure the research question, considering the study outcomes: P: population, I: intervention, C: control, and O: outcomes [12]. The primary research question addressed in this study was: Does the dimensional remodeling of postextraction alveolar defects differ between secondary and primary intention healing?

P (population): Individuals undergoing socket augmentation procedures following tooth extraction.

I (intervention): Socket healing through primary intention following tooth extraction.

C (control): Socket healing through secondary intention following tooth extraction.

O (outcomes): Primary outcomes: changes in socket dimensions during the 3-6 months following tooth extraction. Secondary outcomes: histomorphometric characteristics of bone tissue in sockets healing via primary and secondary intention.

Selection criteria

This systematic review includes randomized controlled trials, observational studies, retrospective analyses, single-blind split-mouth randomized trials, and controlled clinical studies. The primary aim is to evaluate and compare the effects of secondary and primary intention healing on the dimensional changes of the alveolar ridge and the histomorphometric properties of bone tissue following tooth extraction.

Inclusion criteria

- Research articles published within the last decade.
- Full-text studies written in English.
- Human-based research assessing the healing of augmented alveolar defects through secondary and primary intention following tooth extraction.
- Studies analyzing changes in bone dimensions or histomorphometric properties.
- Randomized, prospective, and retrospective research studies.

Exclusion criteria

- Research comparing secondary and primary intention treatments regarding complication rates.
- In vitro and ex vivo studies.
- Studies focusing on only 1 group of sockets with either secondary or primary intention healing.
- Systematic reviews, meta-analyses, case reports, poster and conference presentations, as well as theses.
- Studies involving fewer than 10 patients.

Search strategy

Two independent researchers searched for relevant publications to include in the systematic review. The search was carried out across three databases: Cochrane Library, PubMed, and ScienceDirect. Articles published between March 1, 2013, and March 1, 2023, were considered (Last search date: March 4, 2023). To maintain consistency in keyword combinations across all databases, six Boolean operators ('AND' and 'OR') were used, and keywords were chosen from the MeSH Terms (Medical Subject Headings) thesaurus to ensure the inclusion of

commonly used medical terminology. The preliminary search resulted in the following keyword combination: (((primary intention) AND (secondary intention)) OR ((open flap) AND (closed flap))) AND ((socket) OR (extraction)) AND (tooth).

The article selection process occurred in two phases. In the first phase, duplicate entries were removed, and irrelevant publications were excluded based on titles and abstracts. In the second phase, full-text articles were reviewed, analyzed, and either included in the literature review or rejected according to the defined exclusion and inclusion criteria. Moreover, the references in the selected articles were examined for potentially relevant studies to be included.

Quality assessment

The evaluation of bias risk in prospective randomized studies was conducted using the "RoB 2" tool from The Cochrane Collaboration [13]. This tool includes five established criteria, each with a corresponding evaluation algorithm. Based on the assessment, each criterion is assigned a risk level, classified as low (+), medium (-), or high (x).

Results and Discussion

Study selection

In the first phase of the publication search, a total of 996 articles were initially identified using the selected keywords. After filtering out studies older than ten years, excluding systematic reviews and case reports, and removing duplicates (n=153), 350 articles remained. These articles were reviewed based on their titles and abstracts. Following this, 29 articles were selected for further evaluation. In the second phase, these 29 articles were fully analyzed. Several were excluded based on specific rejection criteria: 11 lacked a control group, 2 focused on complications, 8 did not assess alveolar bone height (BH), bone width (BW), or bone histomorphometric parameters, 1 was a duplicate of another article, 1 was written in Chinese, and 1 did not involve socket augmentation in the control group. Ultimately, 24 articles were excluded, leaving 5 studies [7, 14-17] for inclusion in the systematic review. A diagram of the search process is shown in Figure 1.



Figure 1. PRISMA flowchart search process diagram; BH: bone height change, and BW: bone width change

Characteristics of included studies

A total of 5 studies [7, 14-17] were included in the systematic review. All of these were prospective randomized controlled trials, each with two groups assigned randomly: a control group and an

experimental group. The studies evaluated at least 1 of the predetermined criteria: changes in bone height, changes in bone width, or histological markers of viable bone. One research assessed both alterations in bone height and histological markers [7], three studies focused solely on the histological percentage of new

bone formation [14-16], and one research examined changes in bone height, width, and histomorphometric indicators of bone tissue [17]. The results from all the studies are summarized in **Table 1**.

Statistical analysis

The initial plan included conducting both a systematic review and a meta-analysis, incorporating both qualitative and quantitative methods. However, due to significant differences in the data, a meta-analysis could not be completed. In conclusion, the review focused solely on a descriptive analysis of the data, without quantitative evaluation, to identify relevant information for statistical interpretation. The statistical results were presented as mean values along with standard deviations (mean \pm SD).

Risk of bias assessment

The risk of bias in the included studies was evaluated using the Cochrane RoB 2 tool. All studies were determined to have a low risk of bias overall. However, three studies [7, 14, 17] showed a moderate risk of selective reporting bias among the randomized controlled trials. This did not affect the overall low risk of bias assessment. A visual depiction of the risk of bias analysis, generated using the 'Robvis' tool [18], is shown in **Figure 2**.

Risk of bias domains



Figure 2. Risk of bias assessment of included studies in the review; risk of bias summary; B risk of bias graph; symbols: (+) = low risk of bias, (?) = unclear risk of bias, and (-) = high risk of bias

Impact of primary and secondary healing intention on alveolar height changes

2 studies [7, 17] investigated modifications in alveolar bone height following tooth extraction with secondary and primary intention healing. These studies involved a total of 48 sockets, examining a range of tooth types, including incisors, premolars, canines, and molars. Both studies provided data on the overall reduction in alveolar height post-extraction [7, 17].

In the 2019 research by Aladmawy *et al.* [7], a significant reduction in alveolar height was observed

when healing occurred via primary intention, with an average decrease of -8.1 ± 1.9 millimeters after six months (P = 0.05). For secondary intention healing, the decrease in alveolar height ranged from -7.5 ± 1.8 millimeters (P = 0.05) [14] to -0.9 ± 1.5 millimeters (P < 0.05) [17]. In Seo *et al.*'s study [17], after four months, the modification in alveolar height was -1.4 ± 1.2 millimeters with primary intention healing and -0.9 ± 1.5 millimeters with secondary intention healing, though no statistically significant difference was found between the two healing methods (P = 0.349).

Table 1. Summary of results presented in the studies included in the systematic review

|--|

No.	Study	Type of study	Study sample/Number of study areas	Follow up	BH (mm)	BW (mm)	New bone (%)	BH (mm)	BW (mm)	New bone (%)	Conclusions					
1	Aladmawy <i>et al.</i> [14]	Randomized prospective study	10/20	6 months	-7.5 ± 1.8 (P = 0.389)	-0.1 ± 0.3 (P = 0.317)		-8.1 ± 1.9 (P = 0.389)	0.1 ± 0.5 (P = 0.564)		There were no significant changes observed in bone height (BH) or bone width (BW) with secondary intention healing. In contrast, the primary intention healing group showed a notable increase in the width of keratinized gingiva and a significant reduction in pain levels compared to the secondary intention group.					
7	Aladmawy <i>et al.</i> [15]	Randomized prospective study	8/16	6 months			$71.1\% \pm 23.5\%$ $(P = 0.066)$			$50.9\% \pm 16.2\%$ (P = 0.066)	The study found no statistically important difference in the formation of new bone between primary and secondary intention healing, based on histomorphometric measures.					
3	Barone et al. [16]	Randomized prospective study	34/34	6 months			$22.5\% \pm 3.9\%$ (P = 0.917)			$\begin{array}{l} 22.5\% \pm 4.3\% \\ (\mathrm{P}=0.917) \end{array}$	There was no statistically important difference in histomorphometric parameters between healing via primary intention and secondary intention.					
4	Kim <i>et al.</i> [17]	Randomized prospective study	12/30	6 months			$\begin{array}{l} 47.3\% \ \pm \ 11.3\% \\ (P > 0.05) \end{array}$			$\begin{array}{l} 40.3\% \ \pm 7.8\% \\ (P > 0.05) \end{array}$	Healing of alveolar sockets through secondary intention showed comparable radiological, clinical, and histological results to those observed with primary intention healing.					
S.	Seo <i>et al.</i> [18]	Randomized prospective study	28/28	6 months	-1.4 ± 1.2 (P = 0.349)	-4.9 ± 3.1 (P = 0.529)	$26.2 \pm 17.7\%$ (P > 0.05)	-0.9 ± 1.5 (P = 0.349)	-4.2 ± 2.5 (P = 0.529)	$24.6 \pm 18.4\%$ (P > 0.05)	Both secondary and primary intention healing led to the comparable development of new viable bone and similar radiological alterations in alveolar dimensions.					
		Abbrev	viations: Bl	H = b	Abbreviations: BH = bone height change, BW = bone width change, B = buccal, and L = lingual.											

The impact of primary and secondary healing intention on alveolar width changes

Two studies assessed the modifications in alveolar width following tooth extraction, with a total of 48 sockets included in the analysis [7, 17]. One study focused on anterior teeth, including incisor, canine, and premolar sockets [17], while the other analyzed only posterior teeth, ranging from molars to premolars [14]. In Seo *et al.*'s research [17], after four months, the modification in alveolar width with primary intention healing was -4.9 ± 3.1 millimeters and -4.2 ± 2.5 millimeters with secondary intention healing. However, the difference between these two groups

wasn't statistically important (P = 0.529) [18]. In Aladmawy *et al.*'s study [7], which evaluated alveolar width after six months, the change in width was -0.1 \pm 0.3 millimeters for primary intention healing (P = 0.317) and 0.1 \pm 0.5 millimeters for secondary intention healing [7]. There was no significant statistical difference between the two healing methods regarding alveolar width changes (P = 0.564).

The impact of primary and secondary healing intention on bone histomorphometric indicators

In the systematic review, four studies assessed histomorphometric bone indicators 3-6 months posttooth extraction, focusing on the percentage of new

viable bone formed in the socket during both primary and secondary intention healing [14-17].

Aladmawy *et al.* [14] used an allogeneic freeze-dried mineralized bone substitute (MinerOss, BioHorizons, Birmingham, Alabama, USA) to preserve alveolar dimensions. In the primary intention group, the bone granules were covered with a PTFE membrane and the mucoperiosteal flap was fully sutured. In contrast, the secondary intention group had bone granules left uncovered and no flap suturing. The results displayed that, six months after tooth extraction and alveolar augmentation, new bone formation in the primary intention group was $71.1 \pm 23.5\%$, while the secondary intention group had $50.9 \pm 16.2\%$. The difference was not statistically significant (P = 0.066) [14].

Barone *et al.* [15] used a xenogeneic bone substitute (MP3, Osteobiol, Coazze, Italy) to fill the socket and cover it with a collagen membrane. The results displayed that three months after tooth extraction, the percentage of newly formed viable bone was $22.5 \pm 3.9\%$ in the primary intention group and $22.5\% \pm 4.3\%$ in the secondary intention group, with no significant difference between the groups (P = 0.917) [15].

Kim *et al.* [16] conducted a study using synthetic bone substitutes (Osteon II, Genoss, Suwon, South Korea) in both secondary and primary healing intention groups, with each group having a collagen membrane covering the socket. After 6 months, the primary intention group showed a new bone formation of $47.3 \pm 11.3\%$, while the secondary intention group exhibited $40.3 \pm 7.8\%$. No statistically important difference was observed between the groups (P > 0.05) [16].

In a separate research by Seo *et al.* [17], xenogeneic bone substitutes (InterOss, SigmaGraft, Fullerton, USA) were utilized. In the primary healing group, a buccal coronally displaced mucoperiosteal flap was added on top of the collagen membrane, whereas the secondary healing group had open healing with the membrane left exposed. After four months, the new bone formation was $26.2 \pm 17.7\%$ in the primary intention group and $24.6 \pm 18.4\%$ in the secondary intention group, with no important difference between the two (P > 0.05) [17].

This systematic review analyzed and included 5 studies [7, 14-17], aiming to assess the impact of primary versus secondary intention healing on changes in alveolar socket dimensions and histomorphometric bone indicators.

After tooth extraction, socket remodeling is part of the healing process. Seo *et al.* used cone-beam computed tomography (CBCT) to assess vertical bone loss and found that secondary intention healing resulted in less bone loss compared to primary intention healing,

although the difference wasn't statistically significant [17]. A similar conclusion was drawn in Aladmawy *et al.*'s study [7], which reported no significant differences in socket height or width between the secondary and primary intention healing groups.

Regarding horizontal remodeling of the alveolar ridge, Seo et al. [17] observed similar changes in both healing groups, with no significant differences between secondary and primary intention healing. Similarly, Aladmawy et al. noted horizontal resorption of the alveolar ridge. While the primary intention healing group showed less resorption than the secondary intention group, the difference wasn't statistically significant [7]. The type of materials used-xenogenic bone substitute and collagen membrane in Seo et al.'s research [17], and allogenic bone and a non-resorbable d-PTFE membrane in Aladmawy et al.'s study [7]did not affect the amount of vertical or horizontal resorption in the alveolar ridge. Likewise, Zhao et al. [18] reported a tendency for greater horizontal resorption with primary intention healing when using a xenogenic bone substitute and a collagen membrane, but the difference wasn't statistically significant.

Four studies included in this systematic review analyzed histomorphometric bone parameters following tooth extraction [14-17]. These studies used various materials for socket augmentation, including xenogenic, allogenic bone substitutes, and autogenous bone. Additionally, different types of membranes were used to protect the bone granules: 3 studies employed non-resorbable collagen membranes [15-17], while one utilized a non-resorbable PTFE membrane [14]. Across all studies, higher percentages of new bone formation were observed in the healing of the socket by primary intention, though the differences between primary and secondary healing intention groups were not statistically significant.

Recent histomorphometric studies indicate a trend toward increased new viable bone formation in socket healing by primary intention, likely due to better protection of the bone substitute particles. Gabay et al. [19] understood that primary intention healing led to greater new bone formation, while secondary intention healing resulted in more connective tissue than viable bone. However, in Ramaglia et al.'s case series [20], a higher amount of new bone was observed in secondary intention healing. These findings must be interpreted with caution, as the secondary and primary intention healing groups were assessed at different time points, which may have affected the outcomes.

The evaluation of post-extraction wound healing must extend beyond dimensional and histomorphometric parameters to include clinical indicators such as pain,

swelling, and potential complications like alveolitis, dehiscence, infection, or bleeding. Both primary and secondary healing intentions can involve these issues. Jakse et al. [21] observed that primary intention healing often offers patients a sense of security and comfort, which can accelerate postoperative recovery and lower the risk of socket clot loss. Kilinc and Ataol [22] further supported this by demonstrating that primary intention healing might help reduce the likelihood of alveolitis. In contrast, secondary intention healing minimizes surgical trauma, reduces the need for suturing, and provides greater flexibility in mucosal healing [23]. This approach avoids the need for flap mobilization, preserves the oral vestibule, and retains more keratinized gingiva [7, 23]. Aggarwal et al.'s study [24] reinforced the less traumatic nature of secondary intention healing, reporting that patients in this group experienced less pain and swelling than those in the primary intention group. Rodrigues et al. [25] similarly found significantly reduced pain and swelling in the primary healing group when compared to the secondary intention group.

Conclusion

Based on the results, the method of healing, whether through secondary or primary intention, does not have a significant effect on the dimensional changes observed in the alveolar ridge following tooth extraction. In addition, the type of healing, whether secondary or primary intention, does not affect the formation of new viable bone in the post-extraction site.

Acknowledgments: None

Conflict of Interest: None

Financial Support: None

Ethics Statement: Ethical approval, granted under BEC-OF-57, was obtained from the Bioethical Committee of the Lithuanian University of Health Sciences.

References

- Kim JH, Wadhwa P, Cai H, Kim DH, Zhao BC, Lim HK, et al. Histomorphometric evaluation of socket preservation using autogenous tooth biomaterial and BM-MSC in dogs. Scanning. 2021;2021(10):6676149. doi:10.1155/2021/6676149
- 2. Tadic A, Bajkin B, Mijatov I, Mirnic J, Vukoje K, Sokac M, et al. Influence of L-PRF topical

application on bone tissue healing after surgical extraction of impacted mandibular third molars: Randomized split-mouth clinical study. Appl Sci. 2023;13(8):4823. doi:10.3390/app130848233

- Lin HK, Pan YH, Salamanca E, Lin YT, Chang WJ. Prevention of bone resorption by HA/β-TCP + collagen composite after tooth extraction: A case series. Int J Environ Res Public Health. 2019;16(23):4616. doi:10.3390/ijerph16234616
- Couso-Queiruga E, Stuhr S, Tattan M, Chambrone L, Avila-Ortiz G. Post-extraction dimensional changes: A systematic review and meta-analysis. J Clin Periodontol. 2021;48(1):127-45. doi:10.1111/jcpe.13390
- El-Sioufi I, Oikonomou I, Koletsi D, Bobetsis YA, Madianos PN, Vassilopoulos S. Clinical evaluation of different alveolar ridge preservation techniques after tooth extraction: A randomized clinical trial. Clin Oral Investig. 2023;27(8):4471-80. doi:10.1007/s00784-023-05068-1
- Čandrlić M, Tomas M, Matijević M, Kačarević ŽP, Bićanić M, Udiljak Ž, et al. Regeneration of buccal wall defects after tooth extraction with biphasic calcium phosphate in injectable form vs. Bovine xenograft: A randomized controlled clinical trial. Dent J (Basel). 2023;11(9):223. doi:10.3390/dj11090223
- Aladmawy MA, Natto ZS, Steffensen B, Levi P, Cheung W, Finkelman M, et al. A comparison between primary and secondary flap coverage in ridge preservation procedures: A pilot randomized controlled clinical trial. Biomed Res Int. 2019;2019(7):1-7. doi:10.1155/2019/7679319
- Seyssens L, Eghbali A, Christiaens V, De Bruyckere T, Doornewaard R, Cosyn J. A oneyear prospective study on alveolar ridge preservation using collagen-enriched deproteinized bovine bone mineral and saddle connective tissue graft: A cone beam computed tomography analysis. Clin Implant Dent Relat Res. 2019;21(5):853-61. doi:10.1111/cid.12843
- Kermanshah H, Najafrad E, Valizadeh S. Forced eruption: Alternative treatment approach to restore teeth with subgingival structure. Case Rep Dent. 2022;2022:9521915. doi:10.1155/2022/9521915
- Pasqualini D, Cocero N, Castella A, Mela L, Bracco P. Primary and secondary closure of the surgical wound after removal of impacted mandibular third molars: A comparative study. Int J Oral Maxillofac Surg. 2005;34(1):52-7. doi:10.1016/j.ijom.2004.01.023
- 11. Bello SA, Olaitan AA, Ladeinde AL. A randomized comparison of the effect of partial and

total wound closure techniques on postoperative morbidity after mandibular third molar surgery. J Oral Maxillofac Surg. 2011;69(6):e24-30. doi:10.1016/j.joms.2011.01.025

- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. BMJ. 2021;372:n71. doi:10.1136/bmj.n71
- 13. Flemyng E, Cumpston MS, Arevalo-Rodriguez I, Chandler J, Deeks JJ. Planning a Cochrane review of diagnostic test accuracy. Cochrane Handbook for Systematic Reviews of Diagnostic Test Accuracy. 2023:1-8.
- Aladmawy MA, Natto ZS, Kreitzer M, Ogata Y, Hur Y. Histological and histomorphometric evaluation of alveolar ridge preservation using an allograft and non-resorbable membrane with and without primary closure: A pilot randomized controlled clinical trial. Medicine (Baltimore) 2022;101(26):e29769.

doi:10.1097/MD.000000000029769

- 15. Barone A, Borgia V, Covani U, Ricci M, Piattelli A, Iezzi G. Flap versus flapless procedure for ridge preservation in alveolar extraction sockets: A histological evaluation in a randomized clinical trial. Clin Oral Implants Res. 2015;26(7):806-13. doi:10.1111/clr.12358
- Kim DM, De Angelis N, Camelo M, Nevins ML, Schupbach P, Nevins M. Ridge preservation with and without primary wound closure: A case series. Int J Periodontics Restorative Dent. 2013;33(1):71-8. doi:10.11607/prd.1463
- Seo GJ, Lim HC, Chang DW, Hong JY, Shin SI, Kim G, et al. Primary flap closure in alveolar ridge preservation for periodontally damaged extraction socket: A randomized clinical trial. Clin Implant Dent Relat Res. 2023;25(2):241-51. doi:10.1111/cid.13165
- Zhao LP, Hu WJ, Xu T, Zhan YL, Wei YP, Zhen M, et al. Two procedures for ridge preservation of molar extraction sites affected by severe bone defect due to advanced periodontitis. Beijing Da Xue Xue Bao. 2019;51(3):579-85. doi:10.19723/j.issn.1671-167X.2019.03.03
- Gabay E, Katorza A, Zigdon-Giladi H, Horwitz J, Machtei EE. Histological and dimensional changes of the alveolar ridge following tooth extraction when using collagen matrix and collagen-embedded xenogenic bone substitute: A randomized clinical trial. Clin Implant Dent Relat Res. 2022;24(3):382-90. doi:10.1111/cid.13085

- Ramaglia L, Saviano R, Matarese G, Cassandro F, Williams RC, Isola G. Histologic evaluation of soft and hard tissue healing following alveolar ridge preservation with deproteinized bovine bone mineral covered with xenogenic collagen matrix. Int J Periodontics Restorative Dent. 2018;38(5):737-45. doi:10.11607/prd.3565
- Jakse N, Bankaoglu V, Wimmer G, Eskici A, Pertl C. Primary wound healing after lower third molar surgery: evaluation of 2 different flap designs. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2002;93(1):7-12. doi:10.1067/moe.2002.119519
- 22. Kilinc A, Ataol M. How effective is collagen resorbable membrane placement after partially impacted mandibular third molar surgery on postoperative morbidity? A prospective randomized comparative study. BMC Oral Health. 2017;17(1):126. doi:10.1186/s12903-017-0416-z
- Hong HR, Chen CY, Kim DM, Machtei EE. Ridge preservation procedures revisited: A randomized controlled trial to evaluate dimensional changes with two different surgical protocols. J Periodontol. 2019;90(4):331-8. doi:10.1002/JPER.18-0041
- 24. Aggarwal V, Umarani M, Baliga SD, Issrani R, Prabhu N. Comparison between primary and secondary method of closing surgical wound after tooth extraction: A split-mouth study. Pesqui Bras Odontopediatria Clin Integr. 2021;21(11):e0227. doi:10.1590/pboci.2021.073
- Rodrigues ÉDR, Martins-de-Barros AV, Loureiro AMLC, Carvalho M de V, Vasconcelos B. Comparison of two suture techniques on the inflammatory signs after third molars extraction– A randomized clinical trial. PLoS One. 2023;18(6):e0286413.

doi:10.1371/journal.pone.0286413