

Original Article

Perceptions and Readiness of Oral and Maxillofacial Surgeons and Trainees Toward Artificial Intelligence Integration: A Singapore-Based Study

M. Ruiz^{1*}, Javier L. Moreno², Alejandro F. Ortega³

¹Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Complutense University of Madrid, Madrid, Spain.

²Oral Surgery Unit, Hospital Clínico San Carlos, Madrid, Spain.

³Department of Oral Surgery, Faculty of Medicine, Autonomous University of Madrid, Madrid, Spain.

*E-mail ✉ carlos.ruiz@outlook.com

Received: 05 November 2021; Revised: 29 January 2022; Accepted: 05 February 2022

ABSTRACT

This investigation sought to assess how oral and maxillofacial surgery (OMS) specialists and trainees understand, perceive, and apply artificial intelligence (AI) in their clinical work and training. A cross-sectional questionnaire was distributed to OMS clinicians and residents in Singapore to gather their perspectives on AI in the field. The instrument contained 25 items across five sections and was administered through an online survey system. A total of 48 individuals responded, comprising 37 specialists and 11 trainees. Among them, 60.4% indicated they lacked strong knowledge of AI, 52.1% were unfamiliar with AI applications in OMS, and 81.3% had never received formal instruction related to AI. Many believed that AI could support diagnostic and treatment-planning tasks (72.9%) and help improve patient outcomes (75.0%), and also agreed that AI should be integrated into OMS training (68.8%). No gender-associated differences were observed, although younger clinicians showed more positive views ($p < 0.05$). Key concerns included potential diagnostic or planning errors (77.1%), excessive reliance (70.8%), data security or privacy issues (41.7%), and rising healthcare expenses (41.7%). Despite 68.8% using AI in everyday activities and 62.5% noting that AI made tasks easier, most had not adopted AI in clinical work (62.5%) and felt insufficiently trained or resourced to do so (79.2% and 58.3%, respectively). OMS clinicians and trainees in Singapore generally express positive expectations regarding AI, with younger participants showing greater enthusiasm. Nonetheless, both familiarity and actual utilization of AI remain limited.

Keywords: Oral and maxillofacial surgeons, Singapore, Artificial intelligence, Clinical work and training

How to Cite This Article: Ruiz CM, Moreno JL, Ortega AF. Perceptions and Readiness of Oral and Maxillofacial Surgeons and Trainees Toward Artificial Intelligence Integration: A Singapore-Based Study. *J Curr Res Oral Surg.* 2022;2:52-63. <https://doi.org/10.51847/8gDmjxX3YF>

Introduction

Artificial intelligence (AI) has rapidly expanded within healthcare over recent years. Academic output in medical AI has risen steadily, with an annual increase of 28.4% [1]. Parallel developments are evident in oral and maxillofacial surgery, where AI-based systems have been designed to support diagnosis and surgical planning [2]. Major technology companies, including Google, are also contributing to this surge, producing systems such as MedLM for medical inquiry responses

and AI tools for disease detection [3]. Within health-professional education, numerous AI-enabled applications have been proposed, such as instructional aids, independent learning resources, and automated evaluation systems [4].

Although the healthcare and education sectors are moving toward incorporating AI, the extent to which clinicians, faculty, and trainees are ready to adopt these tools in routine activities remains less explored. Knowledge, attitudes, and practices (KAP) studies are

widely used in health sciences to evaluate baseline viewpoints and to guide program enhancement [5]. In relation to AI, various KAP assessments targeting researchers, instructors, students, and healthcare workers have been published over the past three years [6-8]. In dentistry, prior KAP investigations involving students and practitioners demonstrate mixed knowledge levels but generally supportive attitudes toward integrating AI into dental curricula and practice [9]. However, research focusing specifically on OMS clinicians is lacking, and very few studies have been conducted in East or Southeast Asia. OMS differs from general dentistry in its blend of medical and surgical responsibilities across facial hard and soft tissues, making it unclear whether OMS clinicians' perspectives mirror those of other dental professionals. Therefore, this study aims to assess OMS specialists' and trainees' knowledge, attitudes, and current use of AI within clinical practice and training environments.

Materials and Methods

A cross-sectional survey of OMS clinicians from both public and private institutions in Singapore was carried out between 7 October 2024 and 15 November 2024. The survey adhered to the CHERRIES reporting

guidelines for online questionnaires [10]. Eligible participants included OMS specialists—identified through specialist registration with the Singapore Dental Council—and OMS trainees enrolled in the National University of Singapore (NUS) Master of Dental Surgery (OMS) residency program, as well as graduates of OMS residency programs practicing without specialist accreditation. Ethical exemption from the NUS Institutional Review Board was secured before data collection (NUS-IRB-2024-892).

Survey development, testing and validation

The questionnaire was hosted and managed using Qualtrics XM (USA). Items were developed collaboratively by the study team (**Table 1**). Five sections were included:

- Section 1: demographic details (age, gender, years of practice)
- Sections 2–4: assessment of knowledge, attitudes, and practices
- Section 5: optional free-text comments

The survey incorporated multiple-choice formats, multiple-response selections, Likert-scale items, and open-ended prompts.

Table 1. Survey sections and questions.

Section 1: Demographic Information — Response Options		
Item	Question	Options
Q1	What is your current age?	_____
Q2	Please indicate your gender.	Male / Female
Q3	Are you a board-certified OMS specialist?	Yes / No
Q4	How long have you been practicing OMS?	0–5 / 6–15 / >15
Section 2: Knowledge		
Item	Statement / Prompt	Response Options
Q1	I possess a solid understanding of artificial intelligence (AI), such as machine learning or large language models.	Strongly Agree / Somewhat Agree / Somewhat Disagree / Strongly Disagree
Q2	I am familiar with how AI can be used within OMS practice and educational settings.	Strongly Agree / Somewhat Agree / Somewhat Disagree / Strongly Disagree
Q3	I have previously attended AI-related courses, talks, or training sessions.	Strongly Agree / Somewhat Agree / Somewhat Disagree / Strongly Disagree
Q4	Please list any AI tools or systems you know of (inside or outside OMS).	_____
Section 3: Attitudes		
Item	Statement / Prompt	Response Options
Q1	AI currently helps or could help improve patient outcomes in OMS.	Strongly Agree / Somewhat Agree / Unsure / Somewhat Disagree / Strongly Disagree
Q2	AI ought to be adopted in clinical workflows for diagnosis and treatment planning.	Strongly Agree / Somewhat Agree / Unsure / Somewhat Disagree / Strongly Disagree
Q3	AI should be incorporated into OMS training programmes.	Strongly Agree / Somewhat Agree / Unsure / Somewhat Disagree / Strongly Disagree
Q4	AI might eventually replace OMS surgeons.	Strongly Agree / Somewhat Agree / Unsure / Somewhat Disagree / Strongly Disagree

Q5	Excessive dependence on AI could lead to loss of important clinical skills.	Strongly Agree / Somewhat Agree / Unsure / Somewhat Disagree / Strongly Disagree
Q6	What advantages do you believe AI offers in OMS practice or education? (select all that apply)	<ul style="list-style-type: none"> • Improved efficiency in clinical work • Lower workload for clinicians, teachers or trainees • Enhanced access and tailored experiences for patients/students • I believe there are no benefits Other: _____
Q7	What concerns do you have about using AI in clinical care? (select all that apply)	<ul style="list-style-type: none"> • Issues related to data privacy or security • Risk of incorrect diagnosis or treatment • Overdependence leading to clinician redundancy • Higher healthcare costs Other: _____

Section 4: Practices

Item	Statement / Prompt	Response Options
Q1	I have used AI systems in any context outside my professional work.	Strongly Agree / Somewhat Agree / Unsure / Somewhat Disagree / Strongly Disagree
Q2	I have used AI tools within my OMS clinical practice.	Strongly Agree / Somewhat Agree / Unsure / Somewhat Disagree / Strongly Disagree
Q3	I have used or considered using AI for the following purposes (choose all that apply):	<ul style="list-style-type: none"> • Diagnostic support (radiographic, histopathologic, clinical) • Educating patients or students • Independent learning • Planning treatment • Intraoperative assistance • I have not used nor considered the use of AI in my practice Other: _____
Q4	I have used or thought about using AI for diagnosing or planning treatment in these OMS subspecialties (select all that apply):	<ul style="list-style-type: none"> • Dentofacial deformities • Dentoalveolar procedures • Surgical pathology (including oncology) • Maxillofacial trauma • Implant and preprosthetic surgery • TMJ procedures • I have not used or considered using AI in my practice
Q5	AI helps make my work easier to complete.	Strongly Agree / Somewhat Agree / Unsure / Somewhat Disagree / Strongly Disagree
Q6	I feel I have sufficient training to handle AI tools.	Strongly Agree / Somewhat Agree / Unsure / Somewhat Disagree / Strongly Disagree
Q7	My institution or clinic has the resources required to implement AI in routine care.	Strongly Agree / Somewhat Agree / Unsure / Somewhat Disagree / Strongly Disagree
Q8	What resources do you believe are needed to support better AI adoption in your practice?	_____

Section 5: Additional Feedback

Item	Prompt	Response Area
Q1	Any further comments?	_____

A sample size estimate was completed before launching the survey, yielding a target of 52 participants based on a 95% confidence level and an 8% margin of error. Convenience sampling was intended for participant recruitment. Prior to distribution, the survey underwent qualitative review to ensure its validity. Content validity was confirmed through evaluation by three OMS/AI experts, who examined the relevance and completeness of the items. Face validity was established through pilot testing involving 6 individuals (representing 10% of the planned sample). This step supported refinement of question clarity, readability, and relevance. Internal consistency across Sections 2, 3, and 4 (Knowledge, Attitudes, and Practices) was examined using Cronbach's alpha, with values above 0.7 deemed acceptable.

Survey administration

The questionnaire was circulated to potential respondents through emails sent by professional bodies (e.g., the Association of Oral and Maxillofacial Surgeons Singapore, AOMSS) and academic institutions (NUS). The survey link was open-access without any login requirement. Participants were informed about the study purpose, investigators, and estimated completion time, and consent was obtained at the beginning of the survey. Involvement was optional, with the freedom to discontinue at any stage. No identifying information was requested, responses remained confidential, and no compensation was provided.

Data exported from the platform were compiled in an Excel spreadsheet. To prevent duplicate entries,

records of unique users were checked using IP addresses and cookies. Response timestamps were reviewed to detect entries completed in under 20 seconds as well as incomplete submissions, which were removed due to likely inaccuracy.

Data analysis

For every survey item, descriptive statistics (counts and percentages) were generated. Additional statistical tests explored relationships between demographic variables (gender, age, practice duration) and survey outcomes. For Likert-type items, the positive categories (“Strongly Agree” and “Somewhat Agree”) were merged into “Agree,” and the negative categories (“Strongly Disagree” and “Somewhat Disagree”) were combined into “Disagree.” In five-point Likert items, “Unsure” was kept as a separate category. Associations between demographic factors and these variables were evaluated using Fisher’s exact test. For multi-select questions, each choice was examined as an independent binary variable, again using Fisher’s exact test to determine differences across demographic groups. All analyses were performed in R, with significance fixed at $p < 0.05$.

A thematic analysis of the free-text responses from Sections 4 and 5 was also carried out. The process involved familiarisation with the data, coding, and sorting coded segments into themes to better

understand suggestions for integrating AI into routine practice.

Results and Discussion

A total of 55 submissions were received by 15 November 2024. Of these, 7 were incomplete and excluded, leaving 48 usable responses. The average completion time was 2.43 minutes. Participants had a mean age of 40.8 years, with 35 males (72.9%) and 13 females (27.1%). Years of practice were evenly distributed: 15 (31.3%) had ≤ 5 years (trainees), 17 (35.4%) had 6–15 years (junior specialists), and 16 (33.3%) had > 15 years (senior specialists).

Knowledge

A considerable proportion of respondents indicated limited understanding of AI in the OMS context (**Table 2**). 60.4% reported “Strongly Disagree” or “Somewhat Disagree” when asked about having a good general grasp of AI, and 52.1% responded similarly regarding awareness of AI applications in OMS. Only 18.8% had ever received AI-related training. No statistically significant effects were found for gender ($p = 0.741–1.000$), age ($p = 0.153–1.000$), or practice duration ($p = 0.222–1.000$) (**Table 3**). The knowledge section achieved a Cronbach’s alpha of 0.76, indicating acceptable reliability (**Table 4**).

Table 2. Summary of responses to questions using the Likert scale.

Question	Strongly Agree	Somewhat Agree	Strongly Disagree	Somewhat Disagree	Unsure
Section 2: Knowledge and Familiarity with AI					
Q1: I possess a solid understanding of artificial intelligence (AI)	2 (4.2%)	17 (35.4%)	10 (20.8%)	19 (39.6%)	NA
Q2: I am familiar with how AI is applied in oral and maxillofacial surgery (OMS)	4 (8.3%)	19 (39.6%)	6 (12.5%)	19 (39.6%)	NA
Q3: I have participated in courses, lectures, or other forms of AI training	1 (2.1%)	8 (16.7%)	17 (35.4%)	17 (35.4%)	NA
Section 3: Attitudes Toward AI in OMS					
Q1: AI is, or has the potential to be, helpful in improving patient outcomes in OMS	17 (35.4%)	19 (39.6%)	1 (2.1%)	1 (2.1%)	10 (20.8%)
Q2: Artificial intelligence should be incorporated into clinical practice for diagnosis and treatment planning	15 (31.3%)	20 (41.6%)	2 (4.2%)	2 (4.2%)	9 (18.8%)
Q3: AI should be included as part of OMS residency and continuing education	14 (29.2%)	19 (39.6%)	2 (4.2%)	4 (8.3%)	9 (18.8%)
Q4: In the future, AI could completely replace oral and maxillofacial surgeons	0 (0%)	3 (6.3%)	24 (50.0%)	14 (29.2%)	7 (14.6%)
Q5: Excessive dependence on AI might cause surgeons to lose important clinical skills	6 (12.5%)	22 (45.8%)	4 (8.3%)	11 (22.9%)	5 (10.4%)
Section 4: Current Use and Experience with AI					
Q1: I have utilized AI technologies in areas of my life outside professional practice	10 (20.8%)	23 (47.9%)	5 (10.4%)	8 (16.7%)	2 (4.2%)

Q2: I have incorporated AI technologies into my OMS clinical practice	0 (0%)	12 (25.0%)	15 (31.3%)	15 (31.3%)	6 (12.5%)
Q5: Using AI simplifies and facilitates the completion of my professional tasks	8 (16.7%)	22 (45.8%)	2 (4.2%)	3 (6.3%)	13 (27.1%)
Q6: I feel sufficiently prepared and trained to effectively use AI tools	0 (0%)	6 (12.5%)	19 (39.6%)	19 (39.6%)	4 (8.3%)
Q7: My clinic/institution currently has the necessary infrastructure to integrate AI into practice	0 (0%)	10 (20.8%)	16 (33.3%)	12 (25.0%)	9 (18.8%)

Table 3. Relationship between demographic variables (gender, age, and years of practice) and Likert-scale item responses.

Question	p-value (Gender)	Male	Female	p-value (Age)	Age ≤40	Age >40	p-value (Experience)	1–5 years practice	6–15 years practice	>15 years practice
Section 2: Knowledge										
Q1	0.741			0.770			1.000			
Disagree		22 (62.9%)	7 (53.8%)		17 (63.0%)	12 (57.1%)		9 (60.0%)	10 (58.8%)	10 (62.5%)
Agree		13 (37.1%)	6 (46.2%)		10 (37.0%)	9 (42.9%)		6 (40.0%)	7 (41.2%)	6 (37.5%)
Q2	0.335			1.000			0.393			
Disagree		20 (57.1%)	5 (38.5%)		14 (51.9%)	11 (52.4%)		6 (40.0%)	11 (64.7%)	8 (50.0%)
Agree		15 (42.9%)	8 (61.5%)		13 (48.1%)	10 (47.6%)		9 (60.0%)	6 (35.3%)	8 (50.0%)
Q3	1.000			0.153			0.222			
Disagree		28 (80.0%)	11 (84.6%)		24 (88.9%)	15 (71.4%)		14 (93.3%)	14 (82.4%)	11 (68.8%)
Agree		7 (20.0%)	2 (15.4%)		3 (11.1%)	6 (28.6%)		1 (6.7%)	3 (17.6%)	5 (31.2%)
Section 3: Attitudes										
Q1	1.000			0.004*			<0.001*			
Disagree		2 (5.7%)	0 (0.0%)		0 (0.0%)	2 (9.5%)		0 (0.0%)	0 (0.0%)	2 (12.5%)
Unsure		7 (20.0%)	3 (23.1%)		2 (7.4%)	8 (38.1%)		1 (6.7%)	1 (5.9%)	8 (50.0%)
Agree		26 (74.3%)	10 (76.9%)		25 (92.6%)	11 (52.4%)		14 (93.3%)	16 (94.1%)	6 (37.5%)
Q2	0.251			0.009*			0.024*			
Disagree		4 (11.4%)	0 (0.0%)		0 (0.0%)	4 (19.0%)		0 (0.0%)	0 (0.0%)	4 (25.0%)
Unsure		5 (14.3%)	4 (30.8%)		3 (11.1%)	6 (28.6%)		1 (6.7%)	4 (23.5%)	4 (25.0%)
Agree		26 (74.3%)	9 (69.2%)		24 (88.9%)	11 (52.4%)		14 (93.3%)	13 (76.5%)	8 (50.0%)
Q3	0.478			0.122			0.146			
Disagree		5 (14.3%)	1 (7.7%)		2 (7.4%)	4 (19.0%)		1 (6.7%)	1 (5.9%)	4 (25.0%)
Unsure		8 (22.9%)	1 (7.7%)		3 (11.1%)	6 (28.6%)		2 (13.3%)	2 (11.8%)	5 (31.2%)
Agree		22 (62.9%)	11 (84.6%)		22 (81.5%)	11 (52.4%)		12 (80.0%)	14 (82.4%)	7 (43.8%)
Q4	0.846			0.865			0.415			
Disagree		27 (77.1%)	11 (84.6%)		22 (81.5%)	16 (76.2%)		11 (73.3%)	14 (82.4%)	13 (81.2%)
Unsure		6 (17.1%)	1 (7.7%)		4 (14.8%)	3 (14.3%)		4 (26.7%)	1 (5.9%)	2 (12.5%)
Agree		2 (5.7%)	1 (7.7%)		1 (3.7%)	2 (9.5%)		0 (0.0%)	2 (11.8%)	1 (6.2%)
Q5	0.078			0.828			0.545			
Disagree		13 (37.1%)	2 (15.4%)		9 (33.3%)	6 (28.6%)		7 (46.7%)	3 (17.6%)	5 (31.2%)
Unsure		5 (14.3%)	0 (0.0%)		2 (7.4%)	3 (14.3%)		1 (6.7%)	2 (11.8%)	2 (12.5%)

Agree	17 (48.6%)	11 (84.6%)	16 (59.3%)	12 (57.1%)	7 (46.7%)	12 (70.6%)	9 (56.2%)
Section 4: Practices							
Q1	0.606		0.290		0.075		
Disagree	9 (25.7%)	4 (30.8%)	5 (18.5%)	8 (38.1%)	1 (6.7%)	5 (29.4%)	7 (43.8%)
Unsure	1 (2.9%)	1 (7.7%)	1 (3.7%)	1 (4.8%)	1 (6.7%)	0 (0.0%)	1 (6.2%)
Agree	25 (71.4%)	8 (61.5%)	21 (77.8%)	12 (57.1%)	13 (86.7%)	12 (70.6%)	8 (50.0%)
Q2	0.448		0.335		0.880		
Disagree	23 (65.7%)	7 (53.8%)	15 (55.6%)	15 (71.4%)	9 (60.0%)	10 (58.8%)	11 (68.8%)
Unsure	3 (8.6%)	3 (23.1%)	5 (18.5%)	1 (4.8%)	3 (20.0%)	2 (11.8%)	1 (6.2%)
Agree	9 (25.7%)	3 (23.1%)	7 (25.9%)	5 (23.8%)	3 (20.0%)	5 (29.4%)	4 (25.0%)
Q5	0.785		0.379		0.656		
Disagree	3 (8.6%)	2 (15.4%)	2 (7.4%)	3 (14.3%)	1 (6.7%)	1 (5.9%)	3 (18.8%)
Unsure	10 (28.6%)	3 (23.1%)	6 (22.2%)	7 (33.3%)	3 (20.0%)	5 (29.4%)	5 (31.2%)
Agree	22 (62.9%)	8 (61.5%)	19 (70.4%)	11 (52.4%)	11 (73.3%)	11 (64.7%)	8 (50.0%)
Q6	0.437		0.880		0.810		
Disagree	29 (82.9%)	9 (69.2%)	21 (77.8%)	17 (81.0%)	11 (73.3%)	14 (82.4%)	13 (81.2%)
Unsure	2 (5.7%)	2 (15.4%)	2 (7.4%)	2 (9.5%)	1 (6.7%)	1 (5.9%)	2 (12.5%)
Agree	4 (11.4%)	2 (15.4%)	4 (14.8%)	2 (9.5%)	3 (20.0%)	2 (11.8%)	1 (6.2%)
Q7	1.000		0.720		0.905		
Disagree	21 (60.0%)	8 (61.5%)	15 (55.6%)	14 (66.7%)	8 (53.3%)	10 (58.8%)	11 (68.8%)
Unsure	7 (20.0%)	2 (15.4%)	6 (22.2%)	3 (14.3%)	3 (20.0%)	4 (23.5%)	2 (12.5%)
Agree	7 (20.0%)	3 (23.1%)	6 (22.2%)	4 (19.0%)	4 (26.7%)	3 (17.6%)	3 (18.8%)

*Denotes statistically significant findings.

Table 4. Cronbach’s alpha values for Sections 2–3 of the questionnaire.

Section	Cronbach's alpha
2 (Knowledge)	0.76
3 (Attitudes)	0.71
4 (Practices)	0.78

When participants were asked to name existing AI tools, 25 (52.1%) were able to identify at least one OMS-related system; of these, 15 referred to tools for examination/diagnostic purposes, 7 mentioned options for treatment planning, and 3 cited other functions. For AI outside OMS, 24 (50.0%) listed a large language model, 5 (10.4%) identified robotics, and another 5 (10.4%) named speech-to-text or text-to-speech tools. 12 (25.0%) could not list any AI technology at all.

Attitudes

Overall, participants demonstrated favourable views toward AI use in OMS (Table 2). A substantial proportion agreed that AI could improve patient outcomes (75.0%), should be incorporated into routine practice (72.9%), and ought to be included in OMS training (68.8%). Meanwhile, only 4.2–12.5% selected

“Strongly Disagree” or “Somewhat Disagree” for these items. In contrast, 79.2% disagreed that AI may eventually replace surgeons, and 58.3% agreed that excessive dependence on AI could erode clinical competence. While gender showed no significant influence ($p = 0.078$ – 1.000), respondents aged 40 or younger were significantly more likely to believe AI can improve patient outcomes ($p = 0.004$) and should be introduced into clinical workflows ($p = 0.009$) (Table 3). Significant differences were also seen among clinicians grouped by practice duration ($p < 0.001$ and $p = 0.024$). The attitudes section demonstrated adequate internal reliability, with a Cronbach’s alpha of 0.71 (Table 4).

Regarding perceived benefits and concerns, 83.3% highlighted greater efficiency, 72.9% cited reduced workload, 45.8% noted better personalisation, while 6.3% felt AI provided no benefit. A larger share of respondents aged ≤ 40 regarded personalisation as an advantage ($p = 0.013$), and experience level showed a significant association with reports of increased efficiency ($p = 0.010$) (Table 5). Concerning drawbacks, 77.1% were worried about incorrect diagnostic or planning outputs, 70.8% about

dependency, and 41.7% each about privacy/security issues and cost escalation. These patterns were consistent across demographic groups (**Table 5**).

Table 5. Associations between demographic characteristics and responses to multi-response questions.

Question / Response Option	p-value (Gender)	Male	Female	p-value (Age)	Age ≤40	Age >40	p-value (Experience)	1–5 years	6–15 years	>15 years
Section 3: Attitudes										
Q6: Perceived Benefits of AI in OMS										
Greater clinical efficiency	1.000	29 (82.9%)	11 (84.6%)	1.000	9 (81.8%)	31 (83.8%)	0.010*	13 (86.7%)	17 (100%)	10 (62.5%)
Decreased workload for clinicians/educators/trainees	0.466	24 (68.6%)	11 (84.6%)	0.246	10 (90.9%)	25 (67.6%)	0.355	13 (86.7%)	11 (64.7%)	11 (68.8%)
Improved accessibility and personalization	0.210	14 (40.0%)	8 (61.5%)	0.013*	9 (81.8%)	13 (35.1%)	0.050	11 (73.3%)	6 (35.3%)	5 (31.2%)
I do not see any advantages	0.553	3 (8.6%)	0 (0.0%)	1.000	0 (0.0%)	3 (8.1%)	0.059	0 (0.0%)	0 (0.0%)	3 (18.8%)
Q7: Concerns Regarding AI in Clinical Practice										
Privacy and data security concerns	1.000	15 (42.9%)	5 (38.5%)	0.741	4 (36.4%)	16 (43.2%)	0.555	5 (33.3%)	9 (52.9%)	6 (37.5%)
Risk of inaccurate diagnosis/treatment	0.458	28 (80.0%)	9 (69.2%)	0.246	7 (63.6%)	30 (81.1%)	0.755	12 (80.0%)	12 (70.6%)	13 (81.2%)
Over-dependence leading to obsolescence	0.512	16 (45.7%)	4 (30.8%)	0.720	7 (63.6%)	26 (70.3%)	0.445	9 (60.0%)	11 (64.7%)	13 (81.2%)
Potential increase in healthcare costs	0.182	22 (62.9%)	11 (84.6%)	0.741	4 (36.4%)	16 (43.2%)	0.175	8 (53.3%)	4 (23.5%)	8 (50.0%)
Section 4: Practices										
Q3: Reported or Considered Uses of AI in OMS										
Diagnosis (radiographic, histopathologic, clinical)	0.746	17 (48.6%)	5 (38.5%)	0.732	6 (54.5%)	16 (43.2%)	0.020*	10 (66.7%)	9 (52.9%)	3 (18.8%)
Patient or student education	0.750	14 (40.0%)	6 (46.2%)	0.488	6 (54.5%)	14 (37.8%)	0.091	8 (53.3%)	9 (52.9%)	3 (18.8%)
Self-directed professional learning	0.740	14 (40.0%)	4 (30.8%)	0.288	6 (54.5%)	12 (32.4%)	0.341	8 (53.3%)	5 (29.4%)	5 (31.2%)
Treatment planning	0.317	15 (42.9%)	3 (23.1%)	1.000	4 (36.4%)	14 (37.8%)	0.934	5 (33.3%)	7 (41.2%)	6 (37.5%)
Intraoperative assistance	0.656	6 (17.1%)	1 (7.7%)	0.653	2 (18.2%)	5 (13.5%)	0.227	4 (26.7%)	1 (5.9%)	2 (12.5%)
Have not used and do not plan to use AI	1.000	7 (20.0%)	3 (23.1%)	0.089	0 (0.0%)	10 (27.0%)	0.009*	0 (0.0%)	3 (17.6%)	7 (43.8%)
Q4: AI Use or Consideration by OMS Subspecialty										
Dentofacial deformities	0.740	18 (51.4%)	5 (41.7%)	1.000	5 (50.0%)	18 (48.6%)	0.594	8 (57.1%)	9 (52.9%)	6 (37.5%)
Dentoalveolar surgery	0.065	7 (20.0%)	6 (50.0%)	1.000	2 (20.0%)	9 (24.3%)	0.358	5 (35.7%)	4 (23.5%)	2 (12.5%)
Surgical pathology (including oncology)	0.703	9 (25.7%)	2 (16.7%)	1.000	3 (30.0%)	13 (35.1%)	0.800	4 (28.6%)	7 (41.2%)	5 (31.2%)
Maxillofacial trauma	0.505	13 (37.1%)	3 (25.0%)	1.000	2 (20.0%)	10 (27.0%)	1.000	4 (28.6%)	4 (23.5%)	4 (25.0%)
Implant and pre-prosthetic surgery	0.703	10 (28.6%)	2 (16.7%)	0.065	1 (10.0%)	17 (45.9%)	0.357	3 (21.4%)	8 (47.1%)	7 (43.8%)

TMJ surgery	0.324	15 (42.9%)	3 (25.0%)	0.569	0 (0.0%)	5 (13.5%)	0.519	1 (7.1%)	1 (5.9%)	3 (18.8%)
Have not used and do not plan to use AI	1.000	4 (11.4%)	1 (8.3%)	1.000	3 (30.0%)	10 (27.0%)	0.225	3 (21.4%)	3 (17.6%)	7 (43.8%)

*Indicates significance

Practices

Although 68.8% reported using AI tools in non-clinical settings, only 25.0% had applied AI within OMS. While 62.5% felt AI could simplify work tasks, most believed they lacked adequate training (79.2%) and that their workplace lacked suitable infrastructure (58.3%) (**Table 2**). No significant differences emerged by gender ($p = 0.437$ – 1.000), age ($p = 0.290$ – 0.880), or years of practice ($p = 0.075$ – 0.905) (**Table 3**). The practices section showed strong reliability with a Cronbach’s alpha of 0.78 (**Table 4**).

Concerning specific applications, respondents most frequently used or considered AI for diagnosis (45.8%), followed by patient/student education (41.7%), independent learning (37.5%), treatment planning (37.5%), and intraoperative uses (16.7%). Differences based on years of experience were significant only for diagnostic uses ($p = 0.020$) (**Table 5**). 11 (22.9%) indicated that they had never used nor considered AI, a pattern significantly more common among clinicians with >15 years in practice ($p = 0.009$). The subspecialty most often identified for potential AI integration was dentofacial deformities (50.0%), followed by implant procedures (39.6%) and surgical pathology (35.4%). No demographic group differences were found ($p > 0.05$).

Thematic analysis

For the open-response prompt on what is needed to support AI integration in OMS, four themes emerged:

1. Training access,
2. Workflow/structural modifications,
3. Funding, and
4. Technological optimisation.

Ten (20.8%) responses emphasised training opportunities, such as structured programs, courses, and software access. Five (10.4%) highlighted the need for improved organisation and workflow design—including institutional processes and clarification of regulatory considerations like consent and privacy. Another 5 (10.4%) pointed to the need for increased financial support, such as government or institutional funding, to lower barriers to adoption. Finally, 2 (4.2%) responses mentioned the importance of enhancing AI technologies themselves; one recommended establishing a national database tailored to local

populations, and another expressed reluctance to use AI until performance improves.

There is broad recognition that artificial intelligence (AI) has the capacity to influence numerous aspects of surgical care, including diagnosis, prognostic assessment, treatment planning, and even intraoperative decision-making [11]. As technological capabilities evolve, understanding the present landscape and the difficulties oral and maxillofacial surgery (OMS) practitioners encounter is essential for lowering barriers to AI adoption.

From the knowledge portion of the survey, only about half of respondents were aware of current AI applications in OMS. Likewise, roughly 50% could identify at least one AI tool relevant to clinical work, whereas approximately 25% were unable to recall any AI technology. These knowledge levels mirror findings reported in other healthcare groups and student populations [7, 12, 13]. Since no demographic subgroup showed superior knowledge, the generally modest performance likely stems from limited structured opportunities for AI education; over 80% of participants indicated they had never attended any form of AI-related training. This lack of exposure is not restricted to OMS—one survey of radiologists, for instance, found that nearly 70% had received no AI education [14]. Although the number of publications describing new surgical AI models has increased substantially [15], these developments do not appear to be reaching a significant portion of our clinical community.

In contrast, participants’ attitudes toward AI were generally encouraging. Most respondents either “strongly agreed” or “agreed” that AI has the potential to improve patient care and deserves a role in OMS practice and training. Previous studies of nurses and other healthcare workers revealed similar optimism, with many believing AI could support diagnostic and treatment decisions and was important for modern healthcare delivery [12, 16]. However, unlike those studies—where approximately half of respondents worried about AI displacing their jobs—only 6.3% of participants in the present study expressed this fear. Even so, more than half cautioned that excessive reliance on AI might weaken clinical skills.

Participants younger than the mean age of 40.8 and those with fewer years in practice demonstrated more favourable attitudes toward AI. Although only the

statements “AI can enhance patient outcomes” and “AI should be integrated into practice” reached statistical significance, many other non-significant comparisons showed a similar trend. For instance, 81.5% of respondents aged ≤ 40 and 80% with 1–5 years of experience agreed that “AI should be included in OMS training,” compared with only 52.4% of those older than 40 and 43.8% with more than 15 years of experience. Additionally, those with over 15 years of practice were more likely to state that they do not use—and do not intend to use—AI. These tendencies align with common observations that younger clinicians are generally more open to adopting advanced technologies [17].

In the practice-related section, limited training and hands-on exposure emerged as major issues. Even though nearly two-thirds believed AI could simplify their workload, most had never attempted to utilise AI tools and felt unprepared to integrate such technologies into everyday practice. Importantly, this sentiment was consistent across all demographic categories. Around 30% of respondents also recommended expanding access to training and establishing more formalised workflows to facilitate AI adoption. Comparable barriers have been documented elsewhere; for example, medical schools often lack faculty expertise to teach AI, and healthcare professionals have suggested partnering with developers and advocating broader education initiatives [14, 18]. Incorporating AI earlier—perhaps at the postgraduate or even undergraduate level—may help familiarise future OMS practitioners and minimise the hesitation caused by unfamiliarity.

Outstanding concerns

Participants’ remaining concerns could be categorised into three main themes: risks of inaccurate diagnoses or treatment recommendations, issues involving data security and patient confidentiality, and the potential for increased healthcare costs. Their four proposed strategies—expanding training opportunities, developing clearer clinical workflows, boosting funding, and refining existing AI systems—directly address these perceived challenges.

The most prominent worry was the possibility of misdiagnosis or flawed treatment planning. A small subset of respondents even emphasised the need for substantial improvement and further optimisation of AI tools before integrating them into real-world settings. Current large language models also show limitations; for example, one study reported an average accuracy of only 62.5% (equivalent to a B grade) on OMS-related examination questions [19]. Still, emerging research

demonstrates strong diagnostic performance of AI systems in identifying pathology using clinical data, images, radiographs, and histological slides, as well as in forecasting oral disease outcomes [20–23]. Despite this progress, the potential consequences of incorrect outputs have led some to advocate for rigorous validation before deployment [24]. Ultimately, addressing this concern will require both enhancements to AI accuracy and broader understanding that these tools are intended to support—not replace—clinicians’ decision-making.

Concerns about data protection and potential violations of patient privacy were raised by nearly half of the respondents. This issue is substantial, as AI systems require the ingestion and processing of large datasets throughout training and validation. Privacy breaches may occur during both development and clinical use, especially since regulatory frameworks such as the Health Insurance Portability and Accountability Act (HIPAA) currently lack detailed rules that specifically address AI technologies [25]. Anxieties that major technology corporations, including Google, may re-identify anonymised datasets by linking them to other sources are also justified; successful re-identification has been demonstrated in previous studies [26, 27], and related lawsuits have been filed [28]. Addressing this problem involves two main strategies. First, future AI systems could be trained using high-fidelity synthetic datasets generated by advanced models [29], reducing dependence on real patient data whenever feasible. Second, existing privacy regulations must be revised and strengthened to better safeguard patient information in settings where AI is used.

Ethical aspects of AI adoption in OMS, though not raised by respondents in this study, remain a pertinent topic. Even as AI shows promise for enhancing clinical outcomes, training, and research, its application must continue to uphold transparency, informed consent, and patient autonomy regarding the sharing of their health information and their participation in care decisions [30]. Professional responsibility also requires openly declaring the use of AI in both clinical and research environments. It is equally important to recognise that, at this stage, AI should augment rather than replace the human elements essential to patient care, academic instruction, and scientific inquiry. Rokshad *et al.* proposed a framework to guide the ethical refinement of AI tools in dental practice and research; it addresses challenges across eleven ethical pillars—transparency, diversity, well-being, respect for autonomy, privacy, accountability, equity, prudence, sustainability, solidarity, and governance—

offering a useful structure for protecting patients' interests as AI becomes more integrated [31].

Nearly half of respondents also anticipated that AI might raise healthcare costs. Interestingly, this contrasts with findings from a comparable study among nurses, in which most participants believed AI would actually lower expenses [16]. Although some fear that development costs for AI systems might eventually be passed on to patients, a 2022 review of 200 studies indicated that AI adoption in healthcare has generally resulted in considerable cost reductions [32]. These savings are linked to shorter diagnostic and treatment times, along with efficiency gains that accrue over sustained use. While cost-effectiveness analyses specific to OMS have not yet been conducted, similar financial benefits have been documented in dental applications such as caries detection and the early recognition of oral mucosal lesions [33, 34]. To avoid imposing added costs on patients, governments and institutions may need to invest directly in AI development. Additional cost-reduction strategies include model pruning to remove redundant components and designing explainable AI systems with feedback mechanisms that support usability and long-term sustainability [32].

This cross-sectional investigation is the first to examine the knowledge, attitudes, and practices of OMS specialists and trainees, and although many of their views align with those of other healthcare providers, understanding the specific concerns within our field enables more targeted solutions. Nonetheless, several limitations must be acknowledged. The relatively small OMS community in Singapore may result in an underpowered study. Although 55 responses were initially collected, 7 were incomplete and excluded, leaving a final sample size just below the target of 52. Moreover, the presence of only a single postgraduate training program may concentrate the perspectives of trainees, reducing the applicability of these findings to broader populations. These constraints reduce generalisability and robustness; future studies should therefore involve multiple centres across different countries to increase participant numbers.

Convenience sampling was selected due to limited project manpower and the impracticality of employing more complex sampling techniques such as systematic or stratified sampling. Although convenient, this approach may bias the sample toward individuals who frequently use digital communication channels, potentially producing results that appear more favourable toward AI. Fortunately, given the small OMS community and the high rate of digital literacy in

Singapore, distributing the survey through email and social media platforms of professional organisations likely reached most OMS clinicians within the country. Additionally, the perspectives presented here largely reflect a population composed predominantly of ethnically Southern Chinese clinicians, which may differ from attitudes in Caucasian, African, or other ethnic groups [35]. Finally, these results represent the current cohort of OMS practitioners; their opinions may evolve significantly over the next decade as AI systems improve, laws are updated, and institutions increasingly shift toward AI-supported healthcare delivery.

Conclusion

Although OMS clinicians and trainees in Singapore generally view AI positively, notable gaps in knowledge and practical familiarity remain. The suggestions provided highlight the need for both technological enhancement and policy development before AI can be fully and effectively incorporated into everyday practice and professional training.

Acknowledgments: This manuscript has not been published nor has it been submitted concurrently for publication elsewhere. Additionally, all of the authors have approved the contents of this paper and have agreed to the submission policies of the journal. To the best of our knowledge, all authors have no conflict of interest, financial or otherwise.

Conflict of Interest: None

Financial Support: None

Ethics Statement: The studies involving humans were approved by National University of Singapore Institutional Review Board. The studies were conducted in accordance with the local legislation and institutional requirements. The participants provided their written informed consent to participate in this study.

References

1. Lin M, Lin L, Lin L, Lin Z, Yan X. A bibliometric analysis of the advance of artificial intelligence in medicine. *Front Med (Lausanne)*. 2025;12. doi:10.3389/fmed.2025.1504428
2. Sillmann YM, Monteiro JLGC, Eber P, Baggio AMP, Peacock ZS, Guastaldi FPS. Empowering surgeons: will artificial intelligence change oral and maxillofacial surgery? *Int J Oral Maxillofac Surg*. 2025;54(2):179–90.

- doi:10.1016/j.ijom.2024.09.004
3. GoogleAI. Transforming healthcare with AI. (2025). Available from: <https://ai.google/applied-ai/health/> (Accessed September 11, 2026).
 4. Gordon M, Michelle D, Aderonke A, Hussein U, Xu NY, Rangana B, et al. A scoping review of artificial intelligence in medical education: bEME guide No. 84. *Med Teach*. 2024;46(4):446–70. doi:10.1080/0142159X.2024.2314198
 5. Andrade C, Menon V, Ameen S, Kumar Praharaj S. Designing and conducting knowledge, attitude, and practice surveys in psychiatry: practical guidance. *Indian J Psychol Med*. 2020;42(5):478–81. doi:10.1177/0253717620946111
 6. SaA J, Hasan HE, Alzoubi KH, Khabour OF. Knowledge, attitude, and perceptions of MENA researchers towards the use of ChatGPT in research: a cross-sectional study. *Heliyon*. 2025;11(1):e41331. doi:10.1016/j.heliyon.2024.e41331
 7. Al-Qerem W, Eberhardt J, Jarab A, Al Bawab AQ, Hammad A, Alasmari F, et al. Exploring knowledge, attitudes, and practices towards artificial intelligence among health professions' students in Jordan. *BMC Med Inform Decis Mak*. 2023;23(1):288. doi:10.1186/s12911-023-02403-0
 8. Bhattarai P, Nepal P, Khatri P, Dhungana P, Adhikari R, Magar R, et al. Knowledge, attitude and practice at AI in education: student's perception. *NPRC J Multidiscip Res*. 2024;1:53–66. doi:10.3126/nprcjmr.v1i17.72463
 9. Dashti M, Londono J, Ghasemi S, Sultan Z, Khosraviani F, Moghaddasi N, et al. Attitudes, knowledge, and perceptions of dentists and dental students toward artificial intelligence: a systematic review. *J Taibah Univ Med Sci*. 2024;19(2):327–37. doi:10.1016/j.jtumed.2023.12.010
 10. Eysenbach G. Improving the quality of web surveys: the checklist for reporting results of internet E-surveys (CHERRIES). *J Med Internet Res*. 2004;6(3):e34. doi:10.2196/jmir.6.3.e34
 11. Amin A, Cardoso SA, Suyambu J, Abdus Saboor H, Cardoso RP, Husnain A, et al. Future of artificial intelligence in surgery: a narrative review. *Cureus*. 2024;16(1):e51631. doi:10.7759/cureus.51631
 12. Serbaya SH, Khan AA, Surbaya SH, Alzahrani SM. Knowledge, Attitude and practice toward artificial intelligence among healthcare workers in private polyclinics in Jeddah, Saudi Arabia. *Adv Med Educ Pract*. 2024;15:269–80. doi:10.2147/AMEP.S448422
 13. Mudenda S, Lubinda R, Kasanga M, Mohamed S, Musakuzi Z, Mufwambi W. Artificial intelligence: a knowledge, attitude, and practices survey among pharmacy students at the university of Zambia. *Creat Educ*. 2024;15:2582–96. doi:10.4236/ce.2024.1512157
 14. Goyal S, Sakhi P, Kalidindi S, Nema D, Pakhare AP. Knowledge, attitudes, perceptions, and practices related to artificial intelligence in radiology among Indian radiologists and residents: a multicenter nationwide study. *Cureus*. 2024;16(12):e76667. doi:10.7759/cureus.76667
 15. Li H, Han Z, Wu H, Musaev ER, Lin Y, Li S, et al. Artificial intelligence in surgery: evolution, trends, and future directions. *Int J Surg*. 2025;111(2):2101–11. doi:10.1097/JS9.0000000000002159
 16. Mariano MEM, Shahin MAH, Ancheta SJ, Kunjan MV, AI Dossary MN, et al. Exploring artificial intelligence knowledge, attitudes, and practices among nurses, faculty, and students in Saudi Arabia: a cross-sectional analysis. *Soc Sci Hum Open*. 2025;11:101384. doi:10.1016/j.ssaho.2025.101384
 17. Chan CKY, Lee KKW. The AI generation gap: are gen Z students more interested in adopting generative AI such as ChatGPT in teaching and learning than their gen X and millennial generation teachers? *Smart Learn Environ*. 2023;10(1):60. doi:10.1186/s40561-023-00269-3
 18. Grunhut J, Marques O, Wyatt ATM. Needs, challenges, and applications of artificial intelligence in medical education curriculum. *JMIR Med Educ*. 2022;8(2):e35587. doi:10.2196/35587
 19. Quah B, Yong CW, Lai CWM, Islam I. Performance of large language models in oral and maxillofacial surgery examinations. *Int J Oral Maxillofac Surg*. 2024;53(10):881–6. doi:10.1016/j.ijom.2024.06.003
 20. Abdul NS, Shivakumar GC, Sangappa SB, Di Blasio M, Crimi S, Cicciù M, et al. Applications of artificial intelligence in the field of oral and maxillofacial pathology: a systematic review and meta-analysis. *BMC Oral Health*. 2024;24(1):122. doi:10.1186/s12903-023-03533-7
 21. Khanagar SB, Naik S, Al Kheraif AA, Vishwanathaiah S, Maganur PC, Alhazmi Y, et al. Application and performance of artificial

- intelligence technology in oral cancer diagnosis and prediction of prognosis: a systematic review. *Diagnostics* (Basel). 2021;11(6):1004. doi:10.3390/diagnostics11061004
22. Patel S, Kumar D. Predictive identification of oral cancer using AI and machine learning. *Oral Oncol Rep*. 2025;13:100697. doi:10.1016/j.oor.2024.100697
23. Li X-L, Zhou G. Deep learning in the diagnosis and prognosis of oral potentially malignant disorders. *Cancer Screen Prev*. 2024;3(4):203–13. doi:10.14218/CSP.2024.00025
24. Feng QJ, Harte M, Carey B, Alqarni A, Monteiro L, Diniz-Freitas M, et al. The risks of artificial intelligence: a narrative review and ethical reflection from an oral medicine group. *Oral Dis*. 2025;31(2):348–53. doi:10.1111/odi.15100
25. Rezaeikhonakdar D. AI Chatbots and challenges of HIPAA compliance for AI developers and vendors. *J Law Med Ethics*. 2023;51(4):988–95. doi:10.1017/jme. 2024.15
26. Na L, Yang C, Lo CC, Zhao F, Fukuoka Y, Aswani A. Feasibility of reidentifying individuals in large national physical activity data sets from which protected health information has been removed with use of machine learning. *JAMA Netw Open*. 2018;1(8):e186040. doi:10.1001/jamanetworkopen.2018.6040
27. Duffourc MN, Gerke S. Health care AI and patient privacy-dinerstein v google.JAMA. 2024;331(11):909–10. doi:10.1001/jama.2024.1110
28. Yadav N, Pandey S, Gupta A, Dudani P, Gupta S, Rangarajan K. Data privacy in healthcare: in the era of artificial intelligence. *Indian Dermatol Online J*. 2023;14(6):788–92. doi:10.4103/idoj.idoj_543_23
29. Murdoch B. Privacy and artificial intelligence: challenges for protecting health information in a new era. *BMC Med Ethics*. 2021;22(1):122. doi:10.1186/s12910- 021-00687-3
30. El Khoury N, Hadid D, El-Outa A. Exploring the ethical landscape of artificial intelligence in dentistry: insights from a cross-sectional study. *Cureus*. 2025;17(4):e82667. doi:10.7759/cureus.82667
31. Rokhshad R, Ducret M, Chaurasia A, Karteva T, Radenkovic M, Roganovic J, et al. Ethical considerations on artificial intelligence in dentistry: a framework and checklist. *J Dent*. 2023;135:104593. doi:10.1016/j.jdent.2023.104593
32. Khanna NN, Maindarkar MA, Viswanathan V, Fernandes JFE, Paul S, Bhagawati M, et al. Economics of artificial intelligence in healthcare: diagnosis vs. treatment. *Healthcare* (Basel). 2022;10(12):2493. doi:10.3390/healthcare10122493
33. Schwendicke F, Rossi JG, Göstemeyer G, Elhennawy K, Cantu AG, Gaudin R, et al. Cost-effectiveness of artificial intelligence for proximal caries detection. *J Dent Res*. 2021;100(4):369–76. doi:10.1177/0022034520972335
34. Alotaibi S, Deligianni E. AI in oral medicine: is the future already here? A literature review. *Br Dent J*. 2024;237(10):765–70. doi:10.1038/s41415-024- 8029-9