

Original Article

Early-Life Establishment of the Infant Oral Microbiome: A 15-Month Longitudinal Study of Microbial Succession and Mother-to-Infant Transmission Patterns

Fabio L. Rizzi^{1*}, Lucia F. Romano¹, Ahmed S. Farouk¹

¹Department of Oral Surgery, Faculty of Medicine, University of Padua, Padua, Italy.

*E-mail ✉ fabio.rizzi@outlook.com

Received: 17 October 2023; Revised: 29 January 2024; Accepted: 04 February 2024

ABSTRACT

The oral cavity contains one of the most intricate bacterial ecosystems in humans. Yet, the processes through which newborns first obtain these microorganisms are still not well understood. This work tracked shifts in oral microbial communities among healthy infants and assessed how the maternal oral flora contributes to microbial acquisition in early life. We proposed that oral microbial diversity in infants would rise as they age. A total of 116 whole-saliva samples were obtained from 32 healthy infant–mother pairs at postpartum, 9 months, and 15 months. Bacterial DNA was isolated and analyzed using Human Oral Microbe Identification through Next Generation Sequencing (HOMINGS). Microbial diversity within each dyad (alpha diversity) was quantified with the Shannon index. Differences between infant and maternal microbiomes (beta diversity) were evaluated using weighted Bray–Curtis distances in QIIME 1.9.1. Core microbiome profiles were examined via MicrobiomeAnalyst, and linear discriminant analysis with effect size was employed to detect taxa distinguishing the two groups. Sequencing generated 6,870,571 16S rRNA reads from matched saliva samples. Infant and maternal microbial communities differed significantly ($p < 0.001$). Infant salivary microbial diversity increased over time, whereas mothers displayed a largely consistent core microbiome throughout. Neither breastfeeding nor infant sex influenced diversity patterns. Infants exhibited higher proportions of Firmicutes and lower levels of Actinobacteria, Bacteroidetes, Fusobacteria, and Proteobacteria compared with mothers. SparCC analysis revealed ongoing shifts in infants' microbial interaction networks ($p < 0.05$). Findings indicate that infants begin life with an oral microbiota distinct from that of adults. During the first year, microbial composition and diversity undergo substantial change, and by the period preceding age two, infant oral communities may resemble those of their mothers.

Keywords: Oral microbiota, Microbial diversity, Infant–mother dyads, Postpartum, Initial microbial acquisition

How to Cite This Article: Rizzi FL, Romano LF, Farouk AS. Early-Life Establishment of the Infant Oral Microbiome: A 15-Month Longitudinal Study of Microbial Succession and Mother-to-Infant Transmission Patterns. *J Curr Res Oral Surg.* 2024;4:72-85. <https://doi.org/10.51847/NpV9lui2oj>

Introduction

The human mouth hosts one of the richest bacterial communities, with more than 700 identified species [1, 2].

However, the earliest stages of microbial establishment in infants remain insufficiently described, and the

factors shaping oral colonization during the first year of life are still not fully clarified. Additionally, the long-term health implications of early oral microbial formation are not yet well defined.

Although the oral microbiome plays important roles in growth and immune maturation, the timing of initial

bacterial exposure in infants is debated. Under typical conditions, the fetus develops in a sterile environment. Nevertheless, recent molecular studies have detected bacterial DNA in placental and amniotic samples from healthy pregnancies [3]. The placental microbiome appears more similar to oral bacterial communities than to those of the skin, nasal passages, vagina, or gut [4, 5]. Most early colonizers in infants—largely members of the indigenous oral flora—are acquired during birth or soon after [6]. Emerging research suggests that microbial activity in the placenta may be linked with increased likelihood of preterm delivery [7, 8]. Previous investigations report that the newborn oral cavity is rapidly dominated by *Bifidobacterium* species, particularly *Streptococcus*, *Gemella*, *Veillonella*, *Granulicatella*, and *Rothia*, followed by *Haemophilus*, *Actinomyces*, *Porphyromonas*, *Prevotella*, and *Neisseria* shortly after birth [6-10]. These organisms may contribute to early immune protection and maturation [11, 12]. A range of variables—maternal microbiota, delivery mode, feeding practices, dietary exposures, caregiver interactions, and antibiotic use—can influence microbial acquisition [10, 13-18].

This study analyzed changes in oral microbial communities in healthy infants from the postpartum period through 15 months and explored time-related microbial patterns between infants and mothers. Culture-based detection of *Streptococcus mutans* and 16S rRNA HOMINGS sequencing were used. We posited that newborns harbor a distinctive bacterial community at birth, differing from that of their mothers, and that infant oral microbial diversity increases with age while being shaped by maternal oral microbiota.

Materials and Methods

Study population

This exploratory project was carried out in parallel with a broader intervention study aimed at reducing coercive interactions in couples and parent-child relationships to support healthier daily habits [ClinicalTrials.gov ID: NCT03163082] [19]. From that dataset, 32 family units were chosen at random. Each family was tracked from the child's birth through 15 months of age. Data collection involved structured questionnaires addressing family dynamics, infant feeding, and oral health behaviors at postpartum baseline and again at 9 and 15 months. Follow-up visits took place at the Bellevue Hospital Pediatric Clinic, the Gouverneur Hospital Pediatric Clinic, and at New York University College of Dentistry. Recruitment

procedures and family characteristics are documented elsewhere [20].

Ethics statement

Approval for all study activities was granted by the Institutional Review Boards of the New York University School of Medicine, the New York University College of Dentistry (Research Proposal Oversight Committee), and the New York City Health and Hospital Corporation. Written informed consent was obtained from all participating parents, and permission was provided for their infants' involvement.

Saliva streptococcus mutans assessment

At each time point, saliva was obtained from mothers and infants. Mothers first remained quiet for 5 min, rinsed with sterile water, chewed paraffin for 30 s, and then collected stimulated saliva into chilled 50-ml tubes. Infant samples were gathered using sterile cotton swabs rotated along the oral mucosa and dental ridges for 10–30 s until saturated. Each swab was placed into 2 ml of pre-reduced transport fluid [RTF; [21]] in a labeled vial, and the swab tip was snapped off before sealing. Samples were transported on ice to the New York University College of Dentistry microbiology laboratory. To generate reliable colony-forming unit (CFU) counts, 10 serial 10-fold dilutions (10^{-1} to 10^{-9}) were prepared. A 50- μ l aliquot of each dilution was dispensed onto mitis salivarius agar supplemented with potassium tellurite and bacitracin (MSB; Difco Laboratories, Detroit, MI, United States) using an Autoplate Spiral Plating System (Advanced Instruments, Norwood, MA, United States). Plates were incubated anaerobically for 72 h at 37°C under a gas mixture of 85% N₂, 10% CO₂, and 5% H₂, after which *S. mutans* CFUs were quantified.

Bacterial genomic DNA extraction

Genomic DNA was isolated from 1 ml of saliva (mothers) or from infant swab material using a modified Epicenter DNA purification protocol (Madison, WI, United States) following earlier descriptions [22]. Each sample was treated with 10 μ l proteinase K (10 mg/ml in TES buffer: 10 mM Tris-HCl pH 8.0, 1 mM EDTA, 100 mM NaCl), 10 μ l lysozyme (100 mg/ml in TES), and 2 μ l mutanolysin (5,000 U/ml in PBS), then processed using phenol/chloroform/isoamyl alcohol extraction. DNA quantity and purity were evaluated with a NanoDrop 1000 spectrophotometer (Thermo Scientific, Wilmington, DE, United States). Extracts were stored at -20°C until sequencing.

16S rDNA sequencing and data processing

Libraries targeting the V1–V3 region of the 16S gene were prepared for sequencing using the HOMINGS protocol at the Forsyth Institute Sequencing Facility (Cambridge, MA, United States) as adapted from earlier work [23, 24]. Between 10 and 50 ng of purified DNA served as template for PCR using forward primer 341F

(5'-AATGATACGGCGACCACCGAGATCTACACTA TGGTAATTGTCCTACGGGAGGCAGCAG-3') and reverse primer 806R

(5'-CAAGCAGAAGACGGCATAACGAGATNNNNNN NNNNNNAGTCAGTCAGCCGGACTACHVGGGT WTCTAAT-3'). Amplicons were cleaned with AMPure beads, and a 12 pM denatured library containing 20% PhiX spike-in was loaded onto an Illumina MiSeq (San Diego, CA, United States).

Across samples, sequencing generated more than 50,000 reads on average, with mean read lengths near 441 bp. Any dataset containing fewer than 2,000 bp of usable reads or showing evidence of chimeras was removed. Sequences were screened using QIIME 1.91 “split_libraries.py” with default filters, including a minimum read length of 200 and a minimum quality score of 25, and then joined with “join_paired_ends.py” [25]. Taxonomic assignment relied on 660 species-specific probes and an additional 107 genus-level probes from the Human Oral Microbiome Database (HOMD). ProbeSeq software performed exact-sequence matching [23]. Only quality-approved samples were included for open-reference OTU assignment. OTUs were kept for downstream work if at least 20% of values contained ≥ 4 counts and displayed $\geq 10\%$ variability (inter-quartile range). Final OTU tables were converted to relative abundance profiles for each taxonomic rank [23].

Statistical analysis

Comparisons of relative bacterial abundance were carried out across mother–infant groups, sampling times, infant sex, and breastfeeding categories. Alpha diversity was quantified using the Shannon index through the phyloseq interface in the R vegan package [26]. Diversity outcomes were visualized as multi-group boxplots reflecting the study variables. Group differences were assessed with Wilcoxon–Mann–Whitney tests and the Kruskal–Wallis test. Beta diversity was evaluated using non-phylogenetic Bray–Curtis dissimilarities generated in the QIIME workflow [25]. A heatmap was used to display the key genera that contributed to clustering patterns between maternal and infant saliva samples at each visit.

Principal coordinates analysis (PCoA) was applied to the beta diversity matrices to create two-dimensional ordination plots reflecting sample separation by age point, sex, and feeding mode. Differences in microbial community structure were tested through PERMANOVA, with adjustments for sequencing depth.

To illustrate taxonomic differences between groups, relative abundance profiles at phylum, genus, and species resolution were graphed for mothers versus infants and for breastfed versus non-breastfed infants. Species-level comparisons across postpartum, 9-month, and 15-month assessments were conducted using MaAsLin2 [27]. Core microbiome determination was performed with MicrobiomeAnalyst [28, 29], and the outputs are shown as heatmaps including taxa detected in more than 20% of participants with relative abundance greater than 0.01%. In these heatmaps, the y-axis represents taxon prevalence, while the x-axis corresponds to the detection threshold. Core communities of mothers and infants were compared across all three sampling occasions.

Differentially enriched features were identified using linear discriminant analysis (LDA) with effect size estimation (LEfSe) [30], complemented by nonparametric Kruskal–Wallis testing. Features were considered significant when the FDR-corrected p-value was below 0.1 and the Log LDA score exceeded 2.0. LDA plots reported the discriminating taxa on the y-axis and corresponding LDA values on the x-axis; greater LDA scores indicated higher abundance in mothers. To examine relationships among microbial taxa and developmental trends in infants, SparCC correlation networks were generated with a threshold of 0.5 and $p < 0.05$, where correlated taxa were linked graphically.

For *S. mutans* culture data, CFU counts from MSB agar were converted to log₁₀ values. Mean differences between mothers and infants, as well as comparisons by infant sex and visit time, were evaluated with Wilcoxon–Mann–Whitney tests and the Kruskal–Wallis one-way ANOVA. All analyses were performed using Stata version 17.0 (StataCorp LLC, College Station, TX, United States).

All statistical evaluations were two-tailed, with significance set at $p < 0.05$. Benjamini–Hochberg FDR correction was applied, with $q < 0.05$ indicating statistical significance.

Results and Discussion

At the postpartum baseline visit, 32 mother–infant pairs were enrolled. Maternal ages ranged from 18 to

34 years, with a mean of 24.2 ± 4.2 . All infants were full-term newborns; 22 were male and 10 were female. Reported infant racial distribution included: 15 Latino, 5 African American, 3 Asian, and 9 of mixed background. By the 9- and 15-month visits, 13 dyads returned for follow-up. In total, 116 saliva samples were used for culture-based *S. mutans* evaluation, 16S rDNA sequencing, and HOMINGS profiling.

S. mutans colonization

Based on culture results, *S. mutans* was found in 96.8% of maternal saliva samples at baseline and in 100% of samples at both follow-up assessments. Maternal mean log₁₀ CFU values increased slightly from 4.67 ± 1.23 at postpartum to 4.88 ± 0.99 at 15 months. No infants

carried *S. mutans* at birth. Only 2 infants (15.4%) tested positive at 9 months (mean 1.84 ± 0.34 , log₁₀) and again at 15 months (mean 3.47 ± 1.03 , log₁₀) (**Table 1**). Infant colonization patterns were not related to sex, age, or breastfeeding history.

HOMINGS and 16S rDNA sequencing indicated that the 20 most abundant genera or species accounted for roughly 86.0% of sequences in newborn samples and 88.6% in maternal samples. As summarized in **Table 2**, *Streptococcus* was the dominant genus in infants, comprising more than 72.5% of all identified taxa. At the species level (**Table 3**), *S. mutans* represented 0.0088% of newborn salivary reads and appeared alongside *S. sanguinis* and other oral *Streptococcus* species.

Table 1. Colonization of *S. mutans* in mother–infant dyads.

Time Point	Postpartum Visit	9-Month Well-Child Visit	15-Month Well-Child Visit
Number of mother–infant pairs	N = 32	N = 13	N = 13
	Positive (%)	Mean ± SD	Positive (%)
Infant	0 (0%)	0	2 (15.4%)
Biological Mother	31 (96.8%)	4.67 ± 1.23	13 (100%)

Mean values reflect CFU counts from MSB medium converted to log₁₀ scale.

Table 2. Comparison of the percentage of top 20 16S rDNA genus probes present in the saliva of the mother–infant dyads.

Probe ID	Genus / Species	Relative Abundance in Infants (%)	Relative Abundance in Mothers (%)
GP-081	<i>Streptococcus</i> (Genus probe 4)	71.77%	39.58%
RO-03	<i>Rothia mucilaginosa</i>	10.08%	24.26%
PR-14	<i>Prevotella melaninogenica</i>	0.30%	3.62%
GP-110	<i>Granulicatella</i> (Genus probe)	0.38%	3.00%
ST-20	<i>Streptococcus sanguinis</i>	0.11%	1.82%
GE-04	<i>Gemella sanguinis</i>	0.02%	1.64%
GP-126	<i>Streptococcus</i> (Genus probe 1)	0.14%	1.61%
GE-02	<i>Gemella haemolysans</i>	4.43%	1.32%
PR-09	<i>Prevotella histicola</i>	0.06%	1.19%
GP-060	<i>Neisseria</i> (Genus probe 2)	0.07%	1.17%
GP-004	<i>Actinomyces</i> (Genus probe 4)	0.12%	1.15%
GP-073	<i>Rothia</i> (Genus probe)	0.26%	0.99%
HA-05	<i>Haemophilus parainfluenzae</i>	0.19%	0.91%
GP-089	<i>Veillonella</i> (Genus probe 2)	0.44%	0.63%
PO-09	<i>Porphyromonas</i> sp. oral taxon 279	0.14%	0.60%
RO-02	<i>Rothia dentocariosa</i>	0.01%	0.59%

OR-01	Oribacterium sinus	0.01%	0.55%
SO-01	Solobacterium moorei	0.04%	0.47%
FU-10	Fusobacterium periodonticum	0.07%	0.47%
GP-063	Parvimonas (Genus probe)	0.01%	0.43%

Table 3. Comparison of the percentage of top genus and species of Streptococcus 16S rDNA probes present in the saliva of the mother–infant dyads.”

Probe ID	Target Taxon / Probe Description	Relative Abundance in Infants (%)	Relative Abundance in Mothers (%)
GP-081	Streptococcus (Genus probe 4)	71.7651%	39.5773%
ST-20	Streptococcus sanguinis	0.1148%	1.8175%
GP-126	Streptococcus (Genus probe 1)	0.1411%	1.6107%
ST-22	Streptococcus sp. oral taxon 064	0.1317%	0.0785%
ST-26	Streptococcus sp. oral taxon 431	0.1049%	0.0535%
ST-16	Streptococcus parasanguinis II	0.0789%	0.0448%
ST-23	Streptococcus sp. oral taxon 066	0.0707%	0.0385%
GP-128	Streptococcus (Genus probe 3)	0.0411%	0.0269%
ST-27	Streptococcus sp. oral taxon 486	0.0256%	0.0152%
ST-15	Streptococcus mutans	0.0088%	0.0782%
ST-24	Streptococcus sp. oral taxon 068	0.0064%	0.0063%
GP-127	Streptococcus (Genus probe 2)	0.0039%	0.0024%
ST-09	Streptococcus anginosus	0.0036%	0.0986%
ST-10	Streptococcus constellatus	0.0032%	0.1042%
ST-11	Streptococcus cristatus	0.0031%	0.0016%
ST-14	Streptococcus intermedius	0.0023%	0.0634%
ST-21	Streptococcus sobrinus	0.0007%	0.0671%
ST-12	Streptococcus downei	0.0000%	0.0004%
ST-28	Streptococcus sp. oral taxon 487	0.0000%	0.0004%

Comparison of microbiome diversity between mothers and infants

Sequencing of 116 purified bacterial DNA samples on the MiSeq platform generated a total of 6,870,571 reads, with individual fragments averaging 460 bp (spanning 330–591 bp). Samples contained a mean of 65,270 reads (range 120–151,900; median 47,330). Four samples yielding <1,500 bp were removed before analysis. Out of 767 probes, 600 (78.2%) were positive: 484 (63.5%) in infants and 561 (73.1%) in mothers. Of all assigned sequences, 3,819,592 (55.6%) aligned uniquely to genus-level probes, 2,088,913

(30.4%) matched only one species probe, while 960,108 (14.0%) could not be assigned.

Marked distinctions in community structure and taxonomic distribution were observed between maternal and infant saliva (Wilcoxon; $p < 0.001$). Maternal samples consistently exhibited greater microbial richness at postpartum, 9 months, and 15 months, as indicated by the Shannon index (**Figure 1a**; Kruskal–Wallis, $p < 0.001$). Infant alpha diversity rose steadily at the 9- and 15-month visits compared with the postpartum period, whereas maternal diversity showed minimal temporal change. Beta-diversity analyses (Bray–Curtis distances), visualized via PCoA,

demonstrated clear separation between the two groups at every visit (**Figure 1b**). Maternal samples clustered more tightly than infant samples, reflecting lower within-group variation.




Figure 1. (a) Differences in Shannon diversity across the three infant time points are statistically significant (Kruskal–Wallis). (b) Bray–Curtis comparisons show persistent separation between mothers and infants at all visits ($p < 0.001$, Wilcoxon–Mann–Whitney). (c) Relative percent composition of phyla, genera, and species, displaying the 25 most abundant taxa. Symbols (*) and (^) denote taxa with significant temporal shifts within infants and mothers, respectively (adjusted $p < 0.05$).

Comparison of core oral microbiome between mothers and infants

Figure 1c presents relative abundance distributions across taxonomic levels. Infants showed a

predominance of Firmicutes ($p < 0.05$) and substantially reduced relative levels of Actinobacteria, Bacteroidetes, Fusobacteria, and Saccharibacteria compared with mothers ($p < 0.001$). Among the 25

dominant taxa, infants demonstrated significant temporal variation in *Neisseria flavescens*, *Oribacterium sinus*, *Rothia dentocariosa*, *Veillonella dispar*, *Schaalia sp._HMT_172*, *Proteus mirabilis*, *Abiotrophia defectiva*, *Rothia aeria*, *Fusobacterium periodonticum*, *Streptococcus sanguinis*, *Porphyromonas pasteri*, *Granulicatella elegans*, *Haemophilus parainfluenzae*, *Alloprevotella sp._HMT_473*, *Atopobium parvulum*, and *Prevotella melaninogenica*. Only *Prevotella melaninogenica* changed significantly over time in the maternal cohort. Core microbial genera were defined by $\geq 20\%$ prevalence and $\geq 0.01\%$ relative abundance for infants (Figure 2a) and mothers (Figure 2b) at postpartum (a-

1, b-1), 9 months (a-2, b-2), and 15 months (a-3, b-3). Maternal core taxa were largely constant throughout the 15-month observation period, whereas infants showed substantial expansion of their core community with age. Mothers retained a stable set of genera—including *Streptococcus*, *Rothia*, *Prevotella*, *Gemella*, *Veillonella*, *Neisseria*, and *Actinomyces*—at all visits. Infant saliva initially contained relatively few core taxa but became enriched with additional genera such as *Prevotella*, *Neisseria*, *Alloprevotella*, and *Haemophilus* beginning at the 9-month visit. By 15 months, the divergence in community diversity between infants and mothers had narrowed considerably.




Figure 2. Core microbiome profiles for infants (a) and mothers (b), defined by 20% prevalence and 0.01% abundance thresholds. Maternal profiles remained steady, while infant core taxa broadened over time, with *Neisseria*, *Alloprevotella*, and *Haemophilus* becoming more prominent after 9 months.

Distinguishing taxa between mothers and infants

Genera that differed significantly between mothers and their infants were identified using the linear discriminant analysis effect size approach (LEfSe, FDR 0.05, LDA 2.0) and compared at the postpartum visit (Figure 3a), 9-month visit (Figure 3b), and 15-month visit (Figure 3c). Overall, mothers exhibited a greater number of genera with elevated relative

abundance. In early infancy, *Streptococcus* and *Staphylococcus* appeared at higher levels in infants than in mothers, though this pattern was no longer evident by 15 months. Conversely, *Corynebacterium* was initially more abundant in infants immediately after birth but showed higher levels in mothers at 9 months; this difference also disappeared by 15 months (Figure 3c).




Figure 3. Differentially represented oral genera in mothers and infants during early development. Significant genera were identified by LefSe (FDR 0.05, LDA 2.0) with Kruskal–Wallis testing across the postpartum (a), 9-month (b), and 15-month (c) assessments. Mothers consistently carried more genera with higher abundance. Differences in *Corynebacterium* and *Staphylococcus* diminished steadily over time.

Correlation analysis of oral microbial maturation in infants

Longitudinal comparison of microbial diversity showed that infants experienced a marked rise in salivary microbiome diversity after birth (Kruskal–Wallis, $p < 0.001$). SparCC modeling revealed correlation networks among salivary genera (Figure 4), stratified by infant feeding mode. A correlation cutoff of 0.5 and $p < 0.05$ was applied. The structure of

these microbial networks shifted substantially during early development. At the postpartum time point, *Prevotella* was strongly linked with *Leptotrichia*, *Actinomyces*, and *Granulicatella* (Figure 4a). By 9 months, the network expanded to include *Lautropia*, *Schaalia*, *Neisseria*, *Abiotrophia*, *Porphyromonas*, and *Saccharibacteria* (TM7) (Figure 4b). At 15 months, *Lactobacillus* also became integrated into the network (Figure 4c).




Figure 4. Network mapping of infant oral microbiome progression. Networks were constructed using SparCC correlations at the genus level, categorized by feeding type. Nodes represent genera and edges denote correlation coefficients. The results demonstrate that microbial relationships diversify and become increasingly interconnected as infants age. Threshold: $r = 0.5$, $p < 0.05$. We further evaluated the impact of sex and feeding patterns on infant microbial diversity. Differences in alpha diversity (Shannon index) and beta diversity (Bray–Curtis) between breastfed and formula-fed infants at the postpartum time point were not statistically significant. Compositional distinctions also lacked significance.

Stability of the maternal oral microbiome

We assessed the temporal stability of the salivary microbiome in mothers who completed all visits ($N = 8$). The Shannon diversity index indicated lower diversity at postpartum than at 9 months ($p = 0.02$) and 15 months ($p = 0.001$). However, the Chao1 metric showed no significant shifts over the 15-month period ($p > 0.05$), suggesting an overall stable maternal microbiome. Bray–Curtis beta diversity similarly revealed no significant temporal changes. When infant microbiome stability was compared with their mothers', consistent differences between paired samples were observed at all three time points.

The inclusion of *S. mutans* assessment served three aims: (1) to determine the timing of initial *S. mutans* acquisition in a healthy infant cohort, (2) to compare culture-based detection with 16S rRNA sequencing for identifying *S. mutans* in infant saliva, and (3) to characterize inter- and intra-species interactions between *S. mutans* and other early colonizers. We characterized the oral microbiome of 32 healthy infants at birth, 9 months, and 15 months, with particular emphasis on determinants shaping early microbial establishment. *S. mutans* is a major pathogen implicated in dental caries [31, 32], and early colonization markedly increases disease susceptibility [32–34]. As anticipated, conventional plating detected

S. mutans in 96.8–100% of mothers at all time points. Infants were negative at birth but positive in 15.4% of cases at both 9 and 15 months, consistent with earlier findings [34–36]. Some studies report higher colonization rates in children under 14 months [37, 38], while others indicate a window for initial colonization spanning 7–36 months as primary teeth erupt [39–41]. Given that only two infants were *S. mutans*-positive in our sample, mother–infant transmission patterns could not be conclusively evaluated.

Using the HOMINGS assay, *S. mutans* appeared in small quantities alongside *S. sanguinis* and other *Streptococcus* taxa, supporting its role as an early colonizer and potential anchor for a *Streptococcus*-rich polymicrobial community [42]. Core genera in newborn saliva included *Streptococcus*, *Rothia*, *Gemella*, *Veillonella*, *Staphylococcus*, *Prevotella*, and *Granulicatella*, aligning with previous reports [43]. These observations reinforce the ecological plaque hypothesis [44], emphasizing that early microbial diversification may shape community development and influence lifelong oral health.

We extended our analysis to compare infant oral microbial diversity and genus-level composition at postpartum, 9 months, and 15 months with those of their mothers. Overall, mothers consistently exhibited a richer and more varied salivary microbiome than their

infants. Maternal community structure, diversity indices, and dominant taxa remained largely unchanged across the study period. In contrast, infants showed a pronounced age-related rise in alpha diversity. These observations align with findings from Ramadugu *et al.* who reported that infant salivary richness and diversity increase steadily with age, regardless of maternal oral health, educational background, delivery mode, or feeding practices [45]. Although clear beta-diversity differences were present between mothers and infants at each sampling point, we also noted substantial overlap: 85.7% of infant core genera at postpartum, 80.0% at 9 months, and 90.1% at 15 months were also detectable in the mothers' core microbiome. By 9 and 15 months, infants exhibited more interconnected taxa and denser correlation networks, indicating that their salivary microbial ecosystems undergo continuous restructuring. These patterns suggest that newborn oral cavities begin with a distinct microbial signature, and as colonization expands, the divergence between maternal and infant communities gradually narrows. It is plausible that infant microbial composition may approach maternal-like profiles before the end of the second year of life. Ramadugu *et al.* similarly found that infant salivary communities become increasingly adult-like with age [45], although their study could not identify the specific bacterial contributors. Conversely, Ferrett *et al.* provided evidence for maternal–infant microbial transmission using genomic tracing, strain-level metagenomics, and longitudinal analyses [43]. The biological processes responsible for initial microbial acquisition and the gradual establishment of the infant oral community remain largely unresolved. A prevailing assumption is that bacterial diversity rises with age—a trend supported here. During the first month, *Streptococcus* accounted for 74.8% of the infant salivary microbiome, decreasing to 60.9% at 9 months and 50.0% at 15 months. *Rothia*, the second most abundant genus, increased from 10.5% postpartum to 11.7% and 14.5% at the later visits. *Neisseria* expanded dramatically—by 15.8-fold at 9 months and 77-fold at 15 months. Additional genera, including *Alloprevotella*, *Granulicatella*, *Prevotella*, and *Haemophilus*, also rose progressively during the 15-month window. These patterns mirror the trends documented by Ramadugu *et al.* [45]. *Haemophilus* comprises gram-negative bacteria, many associated with infections. *Neisseria* includes both pathogenic and commensal species commonly residing in the oral cavity [46]. Xu *et al.* observed that *Rothia* levels were significantly reduced in caries-affected preschool children, implying a possible protective

association [47]. Uranga *et al.* reported enhanced interactions among *Rothia*, *Streptococcus*, and *Staphylococcus* during responses to *Rothia*-derived enterobactin biosynthesis [48]. Increases in *Rothia*, *Alloprevotella*, and *Haemophilus* have been linked to oral disease and cancer risk [49, 50]. Although these genera readily colonize mucosal surfaces, including the mouth, our results demonstrate that they are detectable early in infant saliva. The biological significance of their initial acquisition and early composition patterns remains an open question.

Linear discriminant analysis effect size (LEfSe) is designed to evaluate group differences in microbial abundance [30]. It quantifies the strength of associations and identifies discriminatory taxa. In our cohort, LEfSe revealed at least 15 genera with significant mother–infant differences based on Kruskal–Wallis testing. Notably, *Staphylococcus* comprised 1.7% of the infant salivary microbiome postpartum but disappeared from the infant core microbiome by 9 months. Divergences between mother–infant pairs were smaller by 15 months. Reports on *Staphylococcus* colonization in infant saliva are scarce. As a key constituent of the skin microbiome, various species are frequently found in infant saliva, skin, blood, and stool samples [51–53]. Environmental exposure and breastfeeding practices are thought to be primary contributors to early colonization. Another notable finding was the declining significance of *Corynebacterium* differences between mothers and infants from postpartum to 15 months; it was also absent from the infant core microbiome at 9 months. How *Staphylococcus* and *Corynebacterium* contribute to early oral ecological dynamics and how they influence long-term health is still uncertain. The present study adds evidence that may help shape future hypotheses and research directions.

Breast milk serves as a major reservoir of bacteria that can be transferred to an infant's mouth. The relationship between breastfeeding and the promotion or suppression of *S. mutans* in infant saliva has remained debated for many years [38, 54–56]. Much of the existing literature has concentrated on links between breastfeeding practices and the risk of early childhood caries, yet far fewer studies have examined how breastfeeding shapes microbial diversity in the infant oral cavity. Holgerson *et al.* reported in a 2013 cross-sectional study that 3-month-old infants who were breastfed showed distinctly different oral microbial profiles compared with formula-fed infants [57]. Another recent investigation found that breastfed infants exhibited lower salivary microbial diversity at

two months, but this difference was no longer present at 12 months [45].

In our cohort of 32 healthy infants, only five were exclusively breastfed through 15 months. We detected compositional and diversity differences between the feeding groups, with breastfed infants showing increased levels of *Streptococcus* and *Veillonella*, whereas formula-fed infants demonstrated higher levels of *Rothia*. However, the differences were not statistically significant, likely due to the small sample size. These observations should therefore be considered preliminary. As accumulating research highlights the dynamic interplay between the oral environment and microbial colonization, larger, well-powered studies are needed to clarify how feeding patterns influence early microbial acquisition and maturation.

Conclusion

The oral microbiome plays a critical role in maintaining human health [58]. Early microbial establishment contributes substantially to immune development and general well-being in infants. In this study, we utilized the MiSeq sequencing platform, the HOMINGS assay, and bioinformatic analyses to characterize the oral microbiota of 32 healthy newborns from the postpartum period through 15 months—a key window for microbial community formation. Our results indicate that:

- (1) *Streptococcus* is the predominant genus in infant saliva, with *S. mutans* appearing at low levels.
- (2) Noticeable differences in microbial composition persist between mothers and their infants throughout the first 15 months.
- (3) Infants show increasing microbial diversity and shifts in core taxa over time, whereas maternal microbiomes remain comparatively stable.
- (4) No significant differences in microbial diversity were observed based on infant sex or feeding method.

Together, these findings contribute to a clearer understanding of how a health-associated oral microbiome is seeded and develops during infancy.

This work had limitations, particularly the modest number of mother–infant pairs. While 32 dyads were enrolled at postpartum, only 13 completed all follow-up visits. A substantial proportion of sequencing reads could not be identified at the species level, restricting deeper analysis of microbial interactions. As a result, the generalizability of our findings is limited. Moreover, detailed information on maternal oral health—such as caries history or periodontal status—

was unavailable. Future studies would benefit from larger cohorts and more comprehensive maternal data to better elucidate the processes governing oral microbiome establishment in infancy and its potential long-term implications for health.

Acknowledgments: None

Conflict of Interest: None

Financial Support: This study was supported by the National Institute of Dental and Craniofacial Research Grants R01 DE019455, R01 DE013937, R01 DE031025, and NSF-CCF-1934962. Data collection was supported by NIDCR Grant 1R34DE022269-01.

Ethics Statement: The study involving human participants was reviewed and approved by the Institutional Review Boards of the New York University School of Medicine, New York University College of Dentistry (Research Proposal Oversight Committee), and the New York City Health and Hospital Corporation (for the Bellevue Hospital Center) for human subjects participating in research activities. All parents provided informed consent and permitted their children to participate in the study. Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

References

1. Paster BJ, Olsen I, Aas JA, Dewhirst FE. The breadth of bacterial diversity in the human periodontal pocket and other oral sites. *Periodontol*. 2000. 2006;42:80-7. doi:10.1111/j.1600-0757.2006.00174.x
2. Dewhirst FE, Chen T, Izard J, Paster BJ, Tanner AC, Yu WH, et al. The human oral microbiome. *J Bacteriol*. 2010;192:5002-17. doi:10.1128/JB.00542-10
3. Perez-Munoz ME, Arrieta MC, Ramer-Tait AE, Walter J. A critical assessment of the “sterile womb” and “in utero colonization” hypotheses. *Microbiome*. 2017;5:48. doi:10.1186/s40168-017-0268-4
4. Aagaard K, Ma J, Antony KM, Ganu R, Petrosino J, Versalovic J. The placenta harbors a unique microbiome. *Sci Transl Med*. 2014;6:237ra65. doi:10.1126/scitranslmed.3008599
5. Gomez-Arango LF, Barrett HL, McIntyre HD, Callaway LK, Morrison M, Nitert MD. Contributions of the maternal oral and gut microbiome to placental microbial colonization.

- Sci Rep. 2017;7:2860. doi:10.1038/s41598-017-03066-4
6. Makino H, Kushiro A, Ishikawa E, Kubota H, Gawad A, Sakai T, et al. Mother-to-infant transmission of intestinal bifidobacterial strains. *PLoS One*. 2013;8:e78331. doi:10.1371/journal.pone.0078331
 7. Blanc V, O'Valle F, Pozo E, Puertas A, Leon R, Mesa F. Oral bacteria in placental tissues. *Oral Dis*. 2015;21:905-12. doi:10.1111/odi.12364
 8. Prince AL, Ma J, Kannan PS, Alvarez M, Gisslen T, Harris RA, et al. Placental membrane microbiome in preterm birth. *Am J Obstet Gynecol*. 2016;214(5):627.e1-627.e16. doi:10.1016/j.ajog.2016.01.193
 9. Lif Holgerson P, Ohman C, Ronnlund A, Johansson I. Maturation of oral microbiota in children. *PLoS One*. 2015;10:e0128534. doi:10.1371/journal.pone.0128534
 10. Lif Holgerson P, Esberg A, Sjodin A, West CE, Johansson I. Development of the saliva microbiome in infants. *Sci Rep*. 2020;10:9629. doi:10.1038/s41598-020-66658-7
 11. Wade WG. The oral microbiome in health and disease. *Pharmacol Res*. 2013;69:137-43. doi:10.1016/j.phrs.2012.11.006
 12. Yao Y, Cai X, Ye Y, Wang F, Chen F, Zheng C. Role of microbiota in infant health. *Front Immunol*. 2021;12:708472. doi:10.3389/fimmu.2021.708472
 13. Li Y, Caufield PW, Dasanayake AP, Wiener HW, Vermund SH. Mode of delivery influences acquisition of *Streptococcus mutans*. *J Dent Res*. 2005;84:806-11. doi:10.1177/154405910508400905
 14. Thakur R, Singh MG, Chaudhary S, Manuja N. Effect of delivery mode and feeding practices on *Streptococcus mutans*. *Int J Paediatr Dent*. 2012;22:197-202. doi:10.1111/j.1365-263X.2011.01176.x
 15. Biagi E, Quercia S, Aceti A, Beghetti I, Rampelli S, Turrone S, et al. Bacterial ecosystem of mother's milk and infant microbiome. *Front Microbiol*. 2017;8:1214. doi:10.3389/fmicb.2017.01214
 16. Timby N, Domellof M, Holgerson PL, West CE, Lonnerdal B, Hernell O, et al. Oral microbiota in infants fed supplemented formula. *PLoS One*. 2017;12:e0169831. doi:10.1371/journal.pone.0169831
 17. Thompson AL, Houck KM, Jahnke JR. Caesarean delivery and infant microbiome development. *Am J Hum Biol*. 2019;31(2):e23219. doi:10.1002/ajhb.23219
 18. Wu TT, Xiao J, Manning S, Saraithong P, Pattanaporn K, Paster BJ, et al. Mode of delivery and microbiome in caries outcomes. *Front Cell Infect Microbiol*. 2022;12:881899. doi:10.3389/fcimb.2022.881899
 19. Smith Slep AM, Heyman RE, Mitnick DM, Lorber MF, Beauchaine TP. Targeting couple and parent-child coercion. *Behav Res Ther*. 2018;101:82-91. doi:10.1016/j.brat.2017.10.003
 20. Heyman RE, Slep AMS, Lorber MF, Mitnick DM, Xu S, Baucom KJW, et al. Couple CARE program trial. *Prev Sci*. 2019;20(5):620-31. doi:10.1007/s11121-018-0961-y
 21. Syed SA, Loesche WJ. Survival of dental plaque flora in transport media. *Appl Microbiol*. 1972;24:638-44. doi:10.1128/am.24.4.638-644.1972
 22. Li Y, Saxena D, Chen Z, Liu G, Abrams WR, Phelan JA, et al. HIV infection and salivary microbial diversity. *J Clin Microbiol*. 2014;52:1400-11. doi:10.1128/JCM.02954-13
 23. Gomes BP, Berber VB, Kokaras AS, Chen T, Paster BJ. Microbiomes of endodontic-periodontal lesions. *J Endod*. 2015;41:1975-84. doi:10.1016/j.joen.2015.08.022
 24. Mougeot JL, Stevens CB, Cotton SL, Morton DS, Krishnan K, Brennan MT, et al. Concordance of HOMIM and HOMINGS technologies. *J Oral Microbiol*. 2016;8:30379. doi:10.3402/jom.v8.30379
 25. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al. QIIME analysis of sequencing data. *Nat Methods*. 2010;7:335-6. doi:10.1038/nmeth.f.303
 26. McMurdie PJ, Holmes S. Phyloseq: R package for microbiome analysis. *PLoS One*. 2013;8:e61217. doi:10.1371/journal.pone.0061217
 27. Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, et al. Multivariable association discovery in meta-omics. *PLoS Comput Biol*. 2021;17:e1009442. doi:10.1371/journal.pcbi.1009442
 28. Dhariwal A, Chong J, Habib S, King IL, Agellon LB, Xia J. MicrobiomeAnalyst tool. *Nucleic Acids Res*. 2017;45:W180-8. doi:10.1093/nar/gkx295
 29. Chong J, Liu P, Zhou G, Xia J. Using MicrobiomeAnalyst. *Nat Protoc*. 2020;15:799-821. doi:10.1038/s41596-019-0264-1
 30. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, et al. Metagenomic biomarker discovery. *Genome Biol*. 2011;12:R60. doi:10.1186/gb-2011-12-6-r60

31. Loesche WJ. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev.* 1986;50:353-80. doi:10.1128/mr.50.4.353-380.1986
32. Berkowitz RJ. Mutans streptococci: acquisition and transmission. *Pediatr Dent.* 2006;28:106-9.
33. Kohler B, Andreen I, Jonsson B. The earlier the colonization by mutans streptococci, the higher the caries prevalence. *Oral Microbiol Immunol.* 1988;3:14-7. doi:10.1111/j.1399-302X.1988.tb00598.x
34. Ruiz-Rodriguez S, Lacavex-Aguilar V, Pierdant-Perez M, Mandeville P, Santos-Diaz M, Garrocho-Rangel A, et al. Colonization levels of *Streptococcus mutans* between mother and infant. *J Clin Pediatr Dent.* 2014;38:197-200. doi:10.17796/jcpd.38.3.9k6613858x2n82j0
35. Tankunnasombut S, Youcharoen K, Wisuttisak W, Vichayanrat S, Tiranathanagul S. Early colonization of mutans streptococci in Thai children. *Pediatr Dent.* 2009;31:47-51.
36. Warren JJ, Weber-Gasparoni K, Marshall TA, Drake DR, Dehkordi-Vakil F, Dawson DV, et al. Longitudinal study of dental caries risk in young children. *Community Dent Oral Epidemiol.* 2009;37:116-22. doi:10.1111/j.1600-0528.2008.00447.x
37. Wan AK, Seow WK, Purdie DM, Bird PS, Walsh LJ, Tudehope DI. Oral colonization of *Streptococcus mutans* in infants. *J Dent Res.* 2001;80:2060-5. doi:10.1177/00220345010800120701
38. Nizar A, Sheikh M, Khan FR, Iqbal NT, Azam SI, Qureshi S, et al. *Streptococcus mutans* carriage in mothers and children. *Clin Exp Dent Res.* 2022;8:1523-32. doi:10.1002/cre2.648
39. Wan AK, Seow WK, Purdie DM, Bird PS, Walsh LJ, Tudehope DI. Longitudinal study of *Streptococcus mutans* colonization after tooth eruption. *J Dent Res.* 2003;82:504-8. doi:10.1177/154405910308200703
40. Law V, Seow WK, Townsend G. Factors influencing oral colonization of mutans streptococci. *Aust Dent J.* 2007;52:93-100. doi:10.1111/j.1834-7819.2007.tb00471.x
41. Ingemansson Hultquist A, Lingstrom P, Bagesund M. Risk factors for early colonization of mutans streptococci. *BMC Oral Health.* 2014;14:147. doi:10.1186/1472-6831-14-147
42. Liu T, Liu J, Liu J, Yang R, Lu X, He X, et al. Interspecies interactions between *S. mutans* and *S. agalactiae*. *Front Cell Infect Microbiol.* 2020;10:344. doi:10.3389/fcimb.2020.00344
43. Ferretti P, Pasolli E, Tett A, Asnicar F, Gorfer V, Fedi S, et al. Mother-to-infant microbial transmission shapes gut microbiome. *Cell Host Microbe.* 2018;24:133-45.e5. doi:10.1016/j.chom.2018.06.005
44. Marsh PD. Are dental diseases ecological catastrophes? *Microbiology.* 2003;149:279-94. doi:10.1099/mic.0.26082-0
45. Ramadugu K, Bhaumik D, Luo T, Gicquelais RE, Lee KH, Stafford EB, et al. Maternal oral health influences infant microbiome. *J Dent Res.* 2021;100:58-65. doi:10.1177/0022034520947665
46. Tonjum T, Van Putten J. Neisseria. In: Cohen J, Powderly WG, Opal SM, editors. *Infectious diseases.* 4th ed. London: Elsevier; 2017. p.1553-65.
47. Xu L, Chen X, Wang Y, Jiang W, Wang S, Ling Z, et al. Salivary microbiota alterations in preschool children. *Front Physiol.* 2018;9:342. doi:10.3389/fphys.2018.00342
48. Uranga CC, Arroyo P Jr, Duggan BM, Gerwick WH, Edlund A. *Rothia mucilaginosa* produces enterobactin. *mSystems.* 2020;5(2):e00161-20. doi:10.1128/mSystems.00161-20
49. Amer A, Galvin S, Healy CM, Moran GP. Microbiome of oral leukoplakia. *Front Microbiol.* 2017;8:2391. doi:10.3389/fmicb.2017.02391
50. Mougeot JC, Beckman MF, Langdon HC, Brennan MT, Bahrani Mougeot F. Oral microbiome in hematological cancers. *J Clin Med.* 2020;9:4068. doi:10.3390/jcm9124068
51. Klevens RM, Morrison MA, Nadle J, Petit S, Gershman K, Ray S, et al. Invasive MRSA infections in the United States. *JAMA.* 2007;298:1763-71. doi:10.1001/jama.298.15.1763
52. Back-Brito GN, El Ackhar VN, Querido SM, Dos Santos SS, Jorge AO, Reis AS, et al. Oral isolates in HIV-positive patients. *Arch Oral Biol.* 2011;56:1041-6. doi:10.1016/j.archoralbio.2011.02.016
53. Soeorg H, Metsvaht T, Eelmae I, Merila M, Treumuth S, Huik K, et al. Breast milk and neonatal colonization with staphylococci. *Pediatr Res.* 2017;82:759-67. doi:10.1038/pr.2017.150
54. Li Y, Wang W, Caufield PW. Fidelity of mutans streptococci transmission and breastfeeding. *Caries Res.* 2000;34:123-32. doi:10.1159/000016579
55. Nunes AM, Alves CM, Borba de Araujo F, Ortiz TM, Ribeiro MR, Silva AA, et al. Breastfeeding and early childhood caries. *Community Dent Oral*

- Epidemiol. 2012;40(6):542-9. doi:10.1111/j.1600-0528.2012.00703.x
56. Duse M, Zicari AM, Berlutti F, Ernesti I, Occasi F, Leonardi L, et al. Growth of *Streptococcus mutans* in different milks. *Int J Immunopathol Pharmacol.* 2014;27:137-41. doi:10.1177/039463201402700119
57. Holgerson PL, Vestman NR, Claesson R, Ohman C, Domellof M, Tanner AC, et al. Oral microbial profile in breast-fed vs formula-fed infants. *J Pediatr Gastroenterol Nutr.* 2013;56:127-36. doi:10.1097/MPG.0b013e31826f2bc6
58. Bourgeois D, Goncalves LS, Lima-Junior JDC, Carrouel F. Editorial: oral microbiome and systemic health. *Front Microbiol.* 2022;13:855668. doi:10.3389/fmicb.2022.855668