

# **Original Article**

Knowledge and Practices in Treating Pediatric Patients with Systemic Conditions in Riyadh's Dental Community

Amirah Nidhal AlMisfer<sup>1</sup>, Hanan Abdullah AlAbbad<sup>1</sup>, Hussah Abdul Aziz AlHudaithy<sup>1</sup>, Njood Hamad AlSultan<sup>1</sup>, Ohood Khalid AlObairi<sup>1</sup>, Shahzeb Hasan Ansari<sup>2</sup>\*

<sup>1</sup>Dental Interns, Riyadh Elm University, Saudi Arabia. <sup>2</sup>Faculty of Preventive Dentistry, Riyadh Elm University, Saudi Arabia.

\*E-mail 🖂 Shahzebhasan@riyadh.edu.sa

Received: 26 January 2021; Revised: 04 May 2021; Accepted: 05 May 2021

# ABSTRACT

It is well known that good dental hygiene is essential for overall health and a higher standard of living. The oral cavity can help raise an early warning since it may be affected by systemic illness and be the source of its symptoms. Using an online survey tool, this longitudinal survey was conducted among dentists and dental students in Riyadh. Participants were approached by Riyadh's dental clinics, hospitals, and institutions, and asked to complete the survey. A total of 381 responses were obtained from the participants, including 28% general practitioners, 72% students/interns, 66% females, and 34% males. Given that most p-values are less than 0.05, the general practitioners exhibit noticeably greater levels of knowledge than the students/intern group. The participants' general level of awareness and knowledge was below average. Students must be taught about the association between uncommon systemic illnesses in children and how they affect their oral health.

Keywords: Pediatric patients, Oral health, Knowledge, Systemic diseases

How to Cite This Article: AlMisfer AN, AlAbbad HA, AlHudaithy HAA, AlSultan NH, AlObairi OK, Ansari SH. Knowledge and Practices in Treating Pediatric Patients with Systemic Conditions in Riyadh's Dental Community. Ann J Dent Med Assist. 2021;1(1):16-23. https://doi.org/10.51847/5DjQg21WPt

### Introduction

Oral health issues are far more common in children with systemic disorders than in their healthy counterparts. For the child suffering from such conditions, negligence or any kind of delay in resolving the issue might result in serious sickness. As a result, the pediatric dentist is essential in preventing, treating, and stabilizing oral and dental issues that can seriously impair a child's health and standard of living before, during, and following treatment [1-3].

The balance between a healthy person and a sick person can be thrown off both locally and systemically by the oral microbiota [4]. In the oral cavity, bacteria, fungi, viruses, and protozoa can all exist alone or in combination. These may have a significant interaction with the host and each other, which might result in either illness or better health. The following are examples of systemic disorders that present with oral symptoms: autoimmune, hematologic, endocrine, and neoplastic processes. Oral abnormalities in children may be linked to viral etiologies like Kawasaki illness or uncommon malignancies like Langerhans cell histiocytosis [5, 6].

It is generally recognized that maintaining oral health is a critical component of overall health and a higher standard of living. The oral cavity may be the source of symptoms of systemic disease and may be impacted by it; therefore, it can help in putting out an early warning. Consequently, it is important that primary care physicians thoroughly examine the oral cavity to make a diagnosis. Patients' dental health is also negatively impacted by these systemic disorders and their treatments, which is why primary care physicians and

dentists should work together more closely. In most cases, teeth and/or gums are affected by severe oral symptoms of systemic illness. The dental staff must thus take great care and safeguards while treating such oral and/or dental symptoms. Additionally, because of the medications a patient may be taking for these conditions, proper care must be provided [7, 8].

As previously said, dentists need to be aware of these signs to treat pediatric patients efficiently. In general, Hong Kong's medical and dentistry students had a positive outlook and an awareness of the need for cooperation between the two professions in raising care and efficacy standards, according to one of the numerous investigations that have been done to gauge awareness [9].

Dentists need to be more aware of their patients' beliefs, preferences, and anxieties to meet their needs [10]. To prepare future dentists for superior practice in pediatric dentistry, their studies must thus include recommendations and procedures [11]. Little ones with asthma produced significantly higher def, DMF rating, and GI mean scores compared to the control groups, with significant asthmatics regularly showing the greatest def and GI rating, Arafa *et al.* [11] determined the relationship between oral health and childhood asthma. A higher GI mean rating was also linked to a significantly higher average salivary calcium level, a lower stimulated salivary flow rate, and a modified salivary pH, according to the salivary study.

Despite its questionable occurrence in systematic evaluations, dental caries is a complex oral disorder that is frequently found in individuals with obesity and diabetes mellitus. Gingivitis, periodontal disease, and dental cavities are related to similar behaviors, such as poor oral hygiene and bad eating habits. Consuming sugary foods and not cleaning your teeth enough might have more detrimental effects on your oral hygiene. Maintaining dental health will stop chronic oral illnesses and improve the outcomes of long-term inflammatory processes. Thus, a multidisciplinary team of medical and dental specialists is required to treat obese and diabetic individuals [12].

# Study hypotheses

In comparison with practicing dentists, dental students have a lesser understanding of oral health care for children with systemic disorders.

### Aims of the research

- To determine the dentists' and dental students' level of knowledge about the oral health of pediatric patients with systemic disorders.
- To assess how much awareness dentistry students

and dentists possess.

### **Materials and Methods**

## Study design

This is a cross-sectional research carried out among dental students and dentists in Riyadh by an online survey.

#### Study sample

Dental universities, hospitals, and clinics in Riyadh were contacted and participants were demanded to fill up the survey.

# Study instrument

The online questionnaire was formed including questions about personal and demographic data followed by questions linked to systemic diseases and their association with oral health in pediatric patients.

### Instrument validity and reliability

A pilot study was performed by sending the survey to 20 participants and the data was entered into SPSS version 22 to specify the reliability using Cronbach's coefficient alpha. The validity of the questionnaire was examined by sending it to experienced researchers in REU and changes were made according to their feedback and comments.

# Statistical analysis

The gathered data was analyzed using SPSS version 22, where descriptive as well as inferential statistics were performed. Comparisons between groups will be made with the value of significance kept under 0.05 using the chi-square test and correlations were done using Spearman's correlation as the data was not normally distributed.

# **Results and Discussion**

We found that the Cronbach's alpha value for the questionnaire's consistency was 811, indicating high reliability. The power of the sample was also computed, and the result was 0.86, which is also extremely excellent (**Table 1**). There were 381 replies from the participants, with 28% being general practitioners, 72% being students or interns, 66% being female, and 34% being male (**Figures 1 and 2**).

The survey questions and replies are displayed in **Table 2**, along with the corresponding percentages. Notably, only 18.2% of participants said they had a good awareness of the connection between oral health and pediatric systemic disorders. Participants in the study concluded that there was a moderate overall

correlation between dental health and different kinds of systemic disorders. The comparison of survey items by gender is shown in **Table 3**, where most differences were not statistically significant. When asked about the relationship between Down syndrome and oral health, significant comparisons were discovered, with females reporting a stronger correlation than males (P-value = .008). In a similar vein, when asked about the relationship between oral health, heart illness, and asthma, females knew more than males (P-values = .049 and .034, respectively).

Although most of the differences between students/interns and general practitioners were statistically significant, the gender-based comparison yielded varying results when analyzed alongside participant qualifications. Most of the P-values are less than 0.05, indicating that general practitioners have substantially greater levels of knowledge than the students/intern group (**Table 4**).

Lastly, the connection between qualification and the degree of linkage among different systemic disorders is displayed in **Table 5**. It was anticipated that there would be a positive correlation between the degree of relationship and certification of systemic disease with oral health, as assessed by Spearman's correlation, with an enormous amount (P-value < 0.05). Cronbach's alpha in this study was 0.811.

Table 1. Samples' power		
Mean	1.88	
Std. deviation	0.72	
Sample size	381	
Alpha	0.05	
Sample mean	1.98	
Standard error of the mean	0.04	
Critical value	1.94	
Beta	0.14	
Power	0.86	



rigure 1. Gender ratio of study participants



Figure 2. Dentistry levels of study participants

AlMisfer et al., Knowledge and Practices in Treating Pediatric Patients with Systemic Conditions in Riyadh's Dental
Community

Ouestions of survey	Response frequencies
Vour knowledge of redistrie systemic diseases	Low. 29.3% Moderate: 52.6%
Tour knowledge of pediatric systemic diseases	High: 17.0%
	L ou: 27.5%
Your understanding of the connection between pediatric systemic diseases	Low. 27.3% Moderate: 54.2%
and oral health	High: 18 2%
	L ow: 34 5%
The link between leukocyte disorders and oral health in children	Moderate: 46.6%
The link between leakeeyte disorders and oral health in emilden	High: 18 9%
<u></u>	Low: 38.4%
The link between nanillon-lefevre syndrome and oral health in children	Moderate: 40.8%
The link between pupilion lefevre syndrome and oral nearth in emiliter	High: 20.8%
	Low: 26.8%
The link between Down's syndrome and oral health in children	Moderate: 44 7%
	High: 28.4%
	Low: 36.1%
The link between hypophosphatasia and oral health in children	Moderate: 45 3%
	High: 18.7%
	Low: 19.7%
The link between diabetes mellitus and oral health in children	Moderate: 47.4%
	High: 32.9%
	Low: 31.8%
The link between hand, foot, and mouth disease and oral health in children	Moderate: 46.3%
	High: 21.8%
	Low: 44.5%
The link between Kawasaki disease and oral health in children	Moderate: 41.3%
	High: 14.2%
	Low: 46.1%
The link between reiter's syndrome and oral health in children	Moderate: 39.7%
	High: 14.2%
	Low: 32.6%
The link between Peutz-Jeghers syndrome and oral health in children	Moderate: 46.1%
	High: 21.3%
	Low: 35.5%
The link between gastroesophageal reflux and oral health in children	Moderate: 43.4%
	High: 21.1%
	Low: 25%
The link between asthma and oral health in children	Moderate: 47.9%
	High: 27.1%
	Low: 31.6%
The link between Cardiac disease and oral health in children	Moderate: 44.7%
	High: 23.7%
	Low: 34.2%
The link between epilepsy and oral health in children	Moderate: 42.9%
	High: 22.9%
	Yes: 86.1%
Do you feel that you require additional knowledge?	No: 13.9%

**Table 2.** Survey questions and their response frequencies by the study participants

**Table 3.** Gender-based comparison of survey responses among study participants

Questions of the survey	Males	Females	P-value
Understanding pediatric systemic diseases	No statistically significant association		.964
Understanding the relationship between pediatric systemic diseases and oral health	No statistically significant association		.788
The impact of leukocyte disorders on oral health in children	No statistically s	ignificant association	.827

.345
.008*
.079
.088
.524
.372
.444
.161
.250
.049*
.034*
.505
.274

\*Statistically significant (Chi-square test)

Table 4. Comparison of survey responses according to participants' qualifications

1 7 1		1	
Questions of survey	Student/Intern	General practitioner	P-value
	Low: 35%	Low: 16%	
Your comprehension of pediatric systemic diseases	Moderate: 51%	Moderate: 57%	.000*
	High: 14%	High: 27%	
Your knowledge of how pediatric systemic diseases are connected to oral health.	Low: 33%	Low: 12%	
	Moderate: 56%	Moderate: 50%	.000*
	High: 11%	High: 38%	
	Low: 38%	Low: 26%	
The relationship between leukocyte disorders and oral health in	Moderate: 48%	Moderate: 42%	.000*
children	High: 14%	High: 32%	
	Low: 46%	Low: 19%	
The relationship between papilion-lefevre syndrome and oral	Moderate: 39%	Moderate: 46%	.000*
health in children	High: 15%	High: 35%	
	Low: 30%	Low: 29%	
The relationship between Down's syndrome and oral health in	Moderate: 47%	Moderate: 40%	.002*
children	High: 24%	High: 41%	
The relationship between hypophosphatasia and oral health in	Low: 41%	Low: 22%	
	Moderate: 45%	Moderate: 46%	.000*
children	High: 13%	High: 32%	
The relationship between diabetes mellitus and oral health in children	No statistically	significant association	.144
	Low: 38%	Low: 16%	
The relationship between hand, foot, and mouth disease and oral health in children	Moderate: 44%	Moderate: 52%	.000*
	High: 18%	High: 31%	
The relationship between Kawasaki disease and oral health in children	Low: 51%	Low: 29%	
	Moderate: 39%	Moderate: 47%	.000*
	High: 10%	High: 25%	
The relationship between reiter's syndrome and oral health in children	Low: 52%	Low: 31%	
	Moderate: 37%	Moderate: 47%	.001*
	High: 11%	High: 22%	
The relationship between Peutz-Jeghers syndrome and oral health	Low: 38%	Low: 19%	.001*

in children	Moderate: 44%	Moderate: 50%	
	High: 18%	High: 30%	
The relationship between gastroesophageal reflux and oral health in children	Low: 39%	Low: 26%	
	Moderate: 44%	Moderate: 43%	.003*
	High: 17%	High: 31%	
The relationship between asthma and oral health in children	No statistically s	ignificant association	.071
The relationship between cardiac disease and oral health in children	Low: 35%	Low: 22%	
	Moderate: 46%	Moderate: 42%	.001*
	High: 19%	High: 36%	
	Low: 38%	Low: 25%	
The relationship between epilepsy and oral health in children	Moderate: 43%	Moderate: 44%	.015*
	High: 20%	High: 31%	
Do you feel that additional knowledge is required?	No statistically significant association		.097

\*Statistically significant (Chi-square test)

Questions of survey	Dentistry level
Vour commences of systemic illuscess in shildren	rs: .202*
Four comprehension of systemic innesses in culturen	p: .000
V	r <sub>s</sub> : .319*
Tour comprehension of the connection between of a nearth and pediatric systemic disorders	p: .000
The link between children's dental health and laukocyte illnesses	rs:184*
	p: .540
Relationship between children's dental health and Panillon I efevre syndrome	rs: .279*
	p: .000
Relationship between children's dental health and Down syndrome	rs: .169*
	p: .000
Relationship between children's oral health and hypophosphatasia	rs: .235*
	p: .026
Relationship between children's oral health and diabetes mellitus	rs: .100
	p: .052
Relationship between children's dental health and hand foot and mouth diseases	rs: .217*
	p: .000
Palationship between children's dental bealth and Kawasaki illness	rs: .228*
Relationship between enhuren's dentar hearth and Rawasaki niness	p: .000
The link between children's dental health and Reiter's syndrome	rs: .195*
	p: .000
The connection between children's dental health and Pautz Jagher's syndrome	rs: .195*
The connection between children's dental health and Teutz-Jegner's syndrome	p: .000
Deletionship between children's dented bealth and costroesenhoosed reflyy	rs: .166*
Relationship between children's dentai health and gastroesophagear renux	p: .001
The link between shildren's dental bealth and asthma	r <sub>s</sub> : .113
The link between children's dental heard and asuma	p: .028
	rs: .183
Relationship between children's dentai health and caldrac disease	p: .000
Polationship between shildren's dental bealth and enilonsy	rs: .148
Relationship between children's dentai health and ephepsy	p: .004
Do you believe you need additional information?	r <sub>s</sub> : .091
	p: .077

Table 5. Relationship between survey responses and dentistry education level

\*Statistically significant (Spearman's correlation)

The current study sought to determine how wellinformed dental professionals, including students, were about the connections between oral health and several systemic disorders. Interestingly, prior research has mostly concentrated on the connection between systemic disorders and overall oral health. Nevertheless, we focused on children, and after checking many reputable databases, we found no prior research of this kind. However, this section of the study will provide a thorough examination of our findings

concerning the body of current literature.

We asked our research participants about numerous systemic disorders in children and their associations with oral health, including heart problems. In terms of overall answers, only 23% identified a strong link between the aforementioned factors. Furthermore, when these replies were examined based on credentials, it was shown that general practitioners had a substantially greater degree of knowledge about this topic, demonstrating that seniority is important while working with these patients. Similar outcomes were found by Pierdant-Pérez *et al.* [13], who assessed dental practitioners' and students' knowledge of how to treat young patients with endocarditis. Senior dental professionals and residents were shown to be more knowledgeable than juniors.

The general level of awareness was low when asked about specific autosomal illnesses and how they relate to oral health. When asked about Down syndrome, a little better response was noted; this might be because the condition is widespread and dentists commonly encounter individuals with it. However, a muchreduced degree of consciousness was seen when questioned about uncommon disorders, including Reiter's syndrome and Papillon-Lefèvre syndrome, which might be the cause of the inadequate knowledge. Increased exposure to these individuals and cases may assist dental practitioners in learning more, which will raise the likelihood that a treatment will be effective [14].

In both adults and children, diabetes mellitus is another illness that has a strong correlation with oral health. Our results showed a positive number when we asked the participants regarding this correlation. Understanding this is crucial given the rising prevalence of DM in Saudi Arabia. According to local research by AlMutairi et al. [15], periodontal infections and caries are among the oral health issues associated with diabetes mellitus that are highly prevalent. The oral health domain scores of Saudi Arabian children with diabetes mellitus were significantly higher than those of the control group in the current study. In all, 82% of the participants in this research had at least one of the following self-reported oral health issues: periodontitis, xerostomia, mouth pain, and foul breath.

# Study limitations

This research might be enhanced in the future if more data can be gathered in the form of a bigger sample size. Our initial goal was not to compare the ratios of the comparative groups, especially general practitioners and students/interns.

### Conclusion

- The participants' general level of awareness and knowledge was below average.
- There was no discernible gender difference.
- Compared to students/interns, general practitioners had noticeably greater levels of expertise.
- Students must be taught about the link between uncommon systemic illnesses in children and how they affect oral health.

Acknowledgments: The authors of this study would like to acknowledge the support and cooperation of the research center of Riyadh Elm University.

Conflict of Interest: None

Financial Support: None

### Ethics Statement: None

### References

- 1. Thikkurissy S, Lal S. Oral health burden in children with systemic diseases. Dent Clin North Am. 2009;53(2):351-7.
- 2. Da Fonseca MA. Dental care of the pediatric cancer patient. Pediatr Dent. 2004;26(1):53-7.
- El Meligy O, Bahannan S, Hassan M, Eltelety S, Kayal R, Qutob A, et al. Oral health status and habits among 6-13 years old children with limited access to dental care in South Jeddah. Int J Pharm Res Allied Sci. 2019;8(3):109-18.
- 4. Alanazi MH, Barnawi NI, Almohaimel SA, Almutairi MA, Alanezi OK, Qureshi LW, et al. Evaluation of dental pulp testing: simple literature review. Arch Pharm Pract. 2019;10(3):37-40.
- Sampaio-Maia B, Caldas IM, Pereira ML, Pérez-Mongiovi D, Araujo R. The oral microbiome in health and its implication in oral and systemic diseases. Adv Appl Microbiol. 2016;97:171-210.
- 6. Gaddey HL. Oral manifestations of systemic disease. Gen Dent. 2017;65(6):23-9.
- Casamassimo PS, Flaitz CM, Hammersmith K, Sangvai S, Kumar A. Recognizing the relationship between disorders in the oral cavity and systemic disease. Pediatr Clin. 2018;65(5):1007-32.
- 8. Da Fonseca MA. Oral and dental care of local and systemic diseases. Pediatr Dent. 2019:66-76.e2.
- 9. Zhang S, Lo EC, Chu CH. Attitude and awareness of medical and dental students towards collaboration between medical and dental practice in Hong Kong. BMC Oral Health. 2015;15(1):1-6.

- Abdul MS, Qayyum MA, Choudhury D, Saha A, Chandak A, Kumari A. Knowledge on pediatric oral surgery procedures amongst newly qualified post graduate students. J Adv Med Dent Sci Res. 2020;8(5):14-7.
- Arafa A, Aldahlawi S, Fathi A. Assessment of the oral health status of asthmatic children. Eur J Dent. 2017;11(3):357.
- Lifshitz F, Casavalle PL, Bordoni N, Rodriguez PN, Friedman SM. Oral health in children with obesity or diabetes mellitus. Pediatr Endocrinol Rev. 2016;14(2):159-67.
- Pierdant-Pérez M, Peñuelas-Solano E, Girón CB, Guzmán-Uribe D, Ramírez-Carrasco A. Endocarditis prophylaxis in dental practice:

evaluation of knowledge among dental students, pediatric dentistry residents, and specialists in pediatric dentistry. Rev Odont Mex. 2019;23(2):22-6.

- Fadel HT, Baghlaf K, Ben Gassem A, Bakeer H, Alsharif AT, Kassim S. Dental students' perceptions before and after attending a centre for children with special needs: a qualitative study on situated learning. Dent J (Basel). 2020;8(3):69.
- 15. AlMutairi FFJ, Pani SC, Alrobaie FM, Ingle NA. Relationship between type-I diabetes mellitus and oral health status and oral health-related quality of life among children of Saudi Arabia. J Family Med Prim Care. 2020;9(2):647-51.