Asian Journal of Periodontics and Orthodontics
2026, Volume 6, Issue 1, Page No: 96-105
Copyright CC BY-NC-SA 4.0
Available online at: www.tsdp.net

ISSN: 3062-3499

Review Article

Machine Learning-Based Prediction of Periodontal Disease Progression: A Conceptual Framework Integrating Clinical, Behavioral, and Radiographic Determinants

Charles-Ignacio Mendoza-Mollocondo[1]*, Ofelia-Marleny Mamani-Luque[1], Jose-Antonio Supo-Gutierrez[1], Wilson-John Mollocondo-Flores[2], Elsa-Anita Condori-Vilca[3], Yenny Condori-Vilca[3], Lucero-Danitza Mamani-Chipana[4]

[1]Universidad Nacional del Altiplano de Puno, Puno, Peru.
[2]Universidad Nacional Micaela Bastidas de Apurimac, Abancay, Peru.
[3]Institución Educativa Walter Peñaloza Ramella, Peru.
[4]Universidad Nacional de San Agustín de Arequipa, Arequipa, Peru.

***E-mail** ✉ cmendoza@unap.edu.pe

Received: 04 January 2026; **Revised:** 01 April 2026; **Accepted:** 05 April 2026

ABSTRACT

Periodontal disease progression remains a major clinical challenge because deterioration is rarely explained by a single biological or behavioral factor. Changes in probing depth, attachment loss, bleeding, plaque control, smoking exposure, systemic health, and radiographic bone loss interact over time to produce heterogeneous trajectories. Conventional risk assessment approaches often simplify this complexity into static or linear categories. A more integrative approach is needed to support earlier identification of patients at risk for progressive destruction. Existing periodontal prediction methods are limited by their dependence on isolated clinical variables or broad risk categories. Although staging and grading systems have improved diagnostic consistency, they do not fully operationalize longitudinal, multimodal, and nonlinear prediction. Behavioral adherence, radiographic architecture, and site-level disease history may modify risk in ways that conventional tools do not capture. These limitations reduce the clinical usefulness of risk estimates when treatment planning requires individualized prognosis. This article proposes a conceptual framework for machine learning-based prediction of periodontal disease progression. The framework integrates clinical, behavioral, systemic, and radiographic determinants into a unified predictive architecture. It is designed to guide future empirical model development rather than to report new patient data. The central objective is to define how heterogeneous periodontal data can be harmonized, modeled, interpreted, and translated into clinical decision support. The framework is derived from a critical synthesis of peer-reviewed literature published between 2017 and 2026 across periodontology, dental radiology, health informatics, and predictive modeling. It identifies three principal predictor domains: clinical inflammatory and attachment measures, behavioral and systemic modifiers, and radiographic indicators of alveolar bone destruction. It also proposes model selection logic, explainability requirements, and workflow integration principles. Two tables summarize the predictor domains and the proposed framework components. The proposed framework provides a structured roadmap for next-generation periodontal risk prediction tools. By combining multimodal data with interpretable machine learning, it aims to support timely intervention, personalized maintenance planning, and improved communication of progression risk. Its value lies in offering a clinically grounded blueprint for future validation studies, software development, and implementation research. The framework emphasizes that prediction should augment, not replace, professional periodontal judgment.

Keywords: Machine learning, Periodontal disease progression, Conceptual framework, Multimodal prediction, Clinical decision support

How to Cite This Article: Mendoza-Mollocondo CI, Mamani-Luque OM, Supo-Gutierrez JA, Mollocondo-Flores WJ, Condori-Vilca EA, Condori-Vilca Y, et al. Machine Learning-Based Prediction of Periodontal Disease Progression: A Conceptual Framework Integrating Clinical, Behavioral, and Radiographic Determinants. Asian J Periodont Orthodont. 2026;6(1):96-105. https://doi.org/10.51847/3BfE5Jkuxd

Introduction

Periodontitis is a chronic inflammatory disease characterized by progressive destruction of periodontal attachment and supporting alveolar bone, with consequences for tooth retention, function, and quality of life. The 2017 classification framework emphasized

© 2026 Asian Journal of Periodontics and Orthodontics

staging and grading because disease severity and anticipated progression must both inform clinical diagnosis and management [1]. Consensus definitions also clarified that periodontitis reflects complex host–microbial interactions rather than a purely plaque-centered disease process [2]. These conceptual advances created a stronger foundation for risk-oriented periodontal care.

Prediction of progression is clinically important because patients with similar baseline severity may follow different longitudinal courses. Systematic evidence indicates that annual attachment loss, bone level change, and tooth loss vary substantially across populations and follow-up periods [3]. Prognosis is further complicated by site-level and tooth-level factors, including furcation involvement, which can strongly influence long-term tooth survival [4]. Therefore, periodontal prediction must move beyond static diagnosis toward dynamic estimation of future disease worsening.

Current risk assessment approaches remain constrained by limited integration of heterogeneous data. Oral hygiene behavior, smoking exposure, compliance with maintenance, and systemic risk are often evaluated separately from clinical periodontal measurements, despite evidence that these factors contribute to disease burden and treatment outcomes [5, 6]. Smoking, in particular, has a dose-related influence on periodontitis and tooth loss, making it a central behavioral determinant rather than a peripheral covariate [7]. A clinically useful framework should therefore treat behavioral risk as a core predictive domain.

Machine learning offers a potential strategy for integrating diverse periodontal determinants into individualized risk estimates. Early applications have shown that deep learning and machine learning can support periodontal diagnosis, radiographic bone-loss detection, and tooth-level prognosis [8-10]. However, model development without a clear conceptual structure risks producing tools that are technically accurate but poorly aligned with clinical workflows. This article therefore proposes an original conceptual framework for multimodal machine learning-based prediction of periodontal disease progression.

Rationale for a multimodal predictive approach

A multimodal predictive approach is justified because periodontal progression is not reducible to a single clinical measurement. Probing depth, clinical attachment level, bleeding on probing, and tooth mobility capture different aspects of inflammatory burden and structural breakdown. The staging and grading system explicitly recognizes that severity, complexity, and progression risk should be interpreted together rather than separately [3]. Machine learning frameworks can operationalize this principle by allowing multiple data streams to contribute simultaneously to risk estimation.

Clinical indicators are necessary but insufficient because their meaning depends on behavioral and systemic context. A patient with moderate attachment loss but persistent smoking, poor plaque control, and irregular maintenance attendance may face a different risk trajectory from a clinically similar patient with high adherence and stable systemic health. Evidence linking oral hygiene practices to periodontitis supports the need to encode behavioral factors in prediction models [5]. Smoking-related risk further illustrates how behavior modifies the probability of future deterioration even after clinical status is measured [6].

Radiographic determinants add a structural dimension that cannot be fully captured by soft-tissue measurements alone. Alveolar bone level, angular defects, furcation involvement, and bone-loss distribution provide information about cumulative destruction and anatomical vulnerability. Studies of deep learning-based periodontal radiograph interpretation show that automated systems can identify bone loss patterns with clinically relevant accuracy [9, 11, 12]. These findings support the inclusion of radiographic variables as a distinct but complementary predictor domain.

The rationale for machine learning is not simply automation but the capacity to model complex relationships among heterogeneous variables. Periodontal progression may involve nonlinear interactions between baseline bone loss, bleeding, smoking exposure, age, systemic disease, and previous tooth-level prognosis. Machine learning studies in periodontology have demonstrated the feasibility of predictive models using clinical, demographic, salivary, and radiographic inputs [13, 14]. A conceptual framework is needed to specify which inputs should be captured and how they should be interpreted.

A multimodal framework also helps avoid the false choice between clinical expertise and computational prediction. Periodontal staging, grading, and maintenance planning already require professional synthesis of multiple signals, but this synthesis is often implicit and variable between clinicians [2, 15]. Machine learning can make this logic more explicit by generating calibrated risk estimates and interpretable feature contributions. The goal is not to replace periodontal judgment but to provide a reproducible decision-support layer.

Clinical, behavioral, and radiographic data domains

The clinical domain should include measures that reflect current inflammatory activity, historical tissue destruction, and tooth-level complexity. Probing depth, clinical attachment level, bleeding on probing, suppuration, mobility, recession, and plaque indices each provide different information about active disease or vulnerability to further loss. Longitudinal evidence on attachment and bone-level change indicates that these variables are meaningful only when interpreted over time [3]. Therefore, the framework treats clinical features as both baseline descriptors and time-updated predictors.

Probing depth and attachment loss should be engineered as structured features rather than entered only as global averages. Site-level maximum probing depth, number of deep sites, distribution of attachment loss, and change from previous visits may capture disease activity more precisely than a single whole-mouth summary. The diagnostic logic of staging and grading supports this distinction because extent, severity, and progression rate are not interchangeable concepts [1]. Machine learning models should therefore preserve clinically meaningful granularity while avoiding excessive noise from poorly standardized measurements.

The behavioral and systemic domain should include smoking status, cumulative smoking exposure, oral hygiene behaviors, maintenance compliance, dental attendance patterns, diabetes status, glycemic control, medication history, and socioeconomic proxies when ethically and legally appropriate [16-19]. Meta-analytic evidence linking oral hygiene with periodontitis supports inclusion of daily self-care as a modifiable predictor [5]. Smoking is especially important because its effect on periodontitis and tooth loss is both biologically plausible and empirically supported [6, 7]. Systemic and behavioral inputs should be handled carefully because they may influence prediction, intervention planning, and patient communication.

The radiographic domain should capture quantitative alveolar bone loss, defect morphology, furcation status, root proximity, crown-root ratio, and radiographic bone density where measurement quality permits. Furcation involvement has prognostic significance for tooth loss, making it an important anatomical variable in tooth-level prediction [4]. Deep learning studies have shown that panoramic and dental radiographs can be used to detect periodontal bone loss and support periodontitis staging [9, 20]. These findings justify radiographic data as a core modality rather than an optional supplement.

Data quality is central to the proposed framework because machine learning models can amplify measurement inconsistencies. Periodontal charting depends on examiner calibration, radiographic interpretation depends on image quality and projection geometry, and behavioral variables depend on patient reporting. Automated or semi-automated radiographic systems may reduce some sources of variation, but they introduce new requirements for segmentation accuracy, validation across imaging devices, and transparent error handling [21, 22]. Feature engineering should therefore be clinically interpretable and auditable.

The three data domains should be harmonized into a longitudinal patient-tooth-site structure that supports both patient-level risk stratification and tooth-level prognosis. Prior machine learning studies predicting periodontal diagnosis, tooth loss, and bone loss demonstrate that structured clinical and radiographic variables can be transformed into clinically useful prediction tasks [23-25]. **Table 1** summarizes the key predictor variables and their expected contributions within the multimodal framework. This organization creates a bridge between periodontal reasoning and computational model development.

Table 1. Predictor Domains and Variables for a Multimodal Periodontal Disease Progression Framework

Predictor domain	Core variables	Feature-engineering logic	Expected contribution to progression prediction
Clinical inflammatory status	Bleeding on probing, suppuration, plaque index, gingival inflammation	Encode as site-level prevalence, tooth-level burden, and change across visits	Captures current inflammatory activity and modifiable disease burden
Clinical structural destruction	Probing depth, clinical attachment level, recession, mobility	Use maximum values, number of affected sites, distributional summaries, and longitudinal change	Represents accumulated tissue destruction and short-term worsening
Tooth-level anatomical complexity	Furcation involvement, root morphology, crown-root ratio, tooth type	Encode tooth-specific risk modifiers and interaction terms with bone loss	Supports tooth-level prognosis and maintenance planning

Behavioral risk	Smoking status, smoking intensity, oral hygiene behavior, dental attendance, maintenance compliance	Convert into categorical, cumulative, and time-updated adherence features	Captures modifiable exposures that influence future instability
Systemic and contextual risk	Diabetes status, glycemic control, medication history, age, relevant socioeconomic indicators	Use clinically justified variables with fairness review and missing-data assessment	Accounts for host susceptibility and contextual determinants of care
Radiographic bone destruction	Alveolar bone level, percentage bone loss, angular defects, horizontal bone loss	Extract quantitative measures from standardized radiographs or validated automated tools	Measures cumulative structural damage and anatomical progression
Radiographic pattern complexity	Furcation radiolucency, bone density, defect morphology, localized versus generalized bone loss	Combine image-derived features with clinical tooth-level features	Improves discrimination between stable damage and progressive vulnerability

Machine learning model selection and logic

Model selection in periodontal progression prediction should begin with the clinical question rather than with algorithmic novelty. If the task is binary classification, such as predicting whether a patient will experience measurable progression within a defined interval, logistic regression with regularization can provide a transparent baseline. If the task involves nonlinear interactions among probing depth, bone loss, smoking exposure, and maintenance adherence, tree-based models may offer stronger flexibility. Systematic comparisons of machine learning algorithms for periodontitis prediction show that model choice should be judged by discrimination, calibration, interpretability, and clinical usefulness together [14].

Logistic regression remains valuable because it provides an interpretable reference model and allows clinicians to understand directionality in predictor effects. However, periodontal risk is rarely linear, and the relationship between disease severity and future tooth loss may be modified by baseline complexity, compliance, furcation involvement, and systemic risk. Random forest and gradient-boosting methods can capture such interactions while maintaining some interpretability through variable-importance and rule-based explanations. Machine learning studies identifying periodontal disease factors and tooth-loss predictors illustrate the usefulness of these models for structured dental datasets [26, 27].

Neural networks and deep learning methods are most appropriate when the framework incorporates radiographic images or high-dimensional multimodal inputs. Convolutional neural networks have already been applied to periodontal bone-loss detection, periodontitis staging, and radiographic diagnosis using panoramic or intraoral images [9, 20, 28]. These methods can extract visual features that may not be easily represented by manually coded variables. Nevertheless, their use should be justified by sufficient data volume, external validation, and explainability procedures.

The proposed modeling pipeline should include data cleaning, standardization, missing-data handling, feature engineering, model training, internal validation, external validation, calibration assessment, and explainability. Prognostic studies of tooth loss and treatment response show that models must be evaluated against clinically relevant outcomes rather than technical accuracy alone [10, 29, 30]. Calibration is especially important because a poorly calibrated model may exaggerate or underestimate risk even when discrimination appears acceptable. Therefore, the framework prioritizes risk reliability as much as risk ranking.

Interpretability should be embedded into the model-development logic from the beginning. Global explanations can identify whether smoking, bone loss, probing depth, or maintenance adherence drives predictions across the population, while local explanations can clarify why an individual patient is categorized as high risk. Reviews of artificial intelligence in periodontology emphasize that black-box performance is insufficient for responsible clinical translation [31, 32]. In this framework, explainability functions as a clinical safety requirement rather than a decorative analytic add-on.

Integration into clinical decision support

A trained periodontal progression model should be implemented as a decision-support tool within routine periodontal workflows rather than as a separate research interface. The most clinically feasible pathway is integration with electronic dental records, periodontal charting systems, radiographic repositories, and recall-management software. Personalized periodontitis risk estimation from nonimage electronic dental records suggests that structured clinical data can be transformed into

actionable risk scores when data capture is consistent [33]. Such integration would allow prediction to occur at baseline examination, reevaluation, and maintenance visits.

The user interface should communicate risk in a format that supports clinical action without overwhelming the clinician. A model output might include patient-level progression probability, tooth-level risk flags, key contributing variables, and recommended review intervals. Prediction of molar loss and tooth-loss phenotype using machine learning demonstrates that tooth-level outputs can be clinically meaningful when they correspond to decisions about maintenance, extraction risk, and targeted intervention [10, 34]. The framework therefore recommends layered outputs, beginning with a concise risk summary and expanding into interpretable details when needed.

Figure 1 outlines the model-development and clinical-deployment logic required for interpretable periodontal progression prediction.



Figure 1. Model Selection, Explainability, Validation, and Clinical Decision-Support Integration for Periodontal Progression Prediction

Risk communication must be patient-centered because behavioral modification is central to periodontal care. If smoking exposure, oral hygiene behavior, or attendance patterns contribute strongly to predicted progression, the model should present these factors as modifiable targets rather than deterministic judgments. Machine learning tools based on questionnaires, demographic data, and salivary biomarkers show that nonclinical information can support screening, but the resulting risk messages must be ethically framed [35, 36]. The decision-support interface should therefore translate prediction into shared planning, not blame.

Human-in-the-loop validation is essential for maintaining clinical accountability [37-39]. Periodontists should be able to accept, question, or override model recommendations when clinical circumstances are not adequately represented in the data. Comparative work between conventional prognostic systems and machine learning models highlights that algorithmic prediction should be evaluated against established periodontal reasoning rather than assumed superior. The framework positions clinicians as interpreters and supervisors of model output, not passive recipients of automated decisions.

Proposed conceptual framework

The proposed framework begins with structured data capture from three coordinated sources: periodontal charting, behavioral and systemic history, and radiographic assessment. Clinical inputs provide measures of inflammatory activity and tissue destruction, behavioral inputs describe modifiable exposures and adherence, and radiographic inputs quantify structural damage and anatomical complexity. Deep learning approaches to alveolar bone-level measurement demonstrate how image-derived features can complement conventional periodontal records [22]. The first design principle is therefore multimodal data fusion with traceable links to clinical meaning.

Figure 2 illustrates the proposed multimodal pathway from periodontal data capture to interpretable clinical action.



Figure 2. Multimodal Conceptual Framework for Machine Learning-Based Prediction of Periodontal Disease Progression

The second stage is data harmonization, which converts heterogeneous inputs into comparable, temporally ordered features. Probing depth, attachment level, bleeding, smoking status, maintenance attendance, and bone-loss measurements should be aligned by patient, tooth, site, and time point. Studies estimating alveolar bone loss and automatically recognizing teeth on radiographs show that automated radiographic features can be integrated with structured clinical variables if measurement definitions are standardized [23, 25]. Harmonization is therefore the technical bridge between periodontal examination and machine learning.

The third stage is model development, where algorithm choice is matched to the prediction task. A transparent model may be preferred for chairside risk communication, whereas a hybrid architecture may be suitable when images and structured records are combined. Evidence from diagnostic deep learning, screening models, and tooth-loss prediction indicates that no single algorithm is universally optimal across periodontal tasks [8, 24, 35]. The framework recommends benchmarking simple and complex models before selecting the final clinical candidate.

The fourth stage is interpretation and clinical translation. The model should generate calibrated progression probabilities, identify influential

predictors, and suggest clinically relevant risk categories. Studies of machine learning-assisted treatment-response prediction and personalized periodontitis risk show that predictive outputs become more useful when connected to care planning, reevaluation intervals, and preventive strategies [30, 33]. **Table 2** presents the core components and design principles of the proposed framework.

Table 2. Proposed Conceptual Framework for Machine Learning-Based Periodontal Progression Prediction: Components, Design Principles, and Integration Logic

Framework component	Design principle	Integration logic	Clinical function
Data capture	Collect clinical, behavioral, systemic, and radiographic inputs using standardized definitions	Link periodontal charting, patient history, and radiographic records within one longitudinal structure	Creates a comprehensive baseline for individualized prognosis
Data harmonization	Align variables by patient, tooth, site, and time	Transform heterogeneous data into structured features suitable for modeling	Reduces fragmentation between clinical examination and radiographic interpretation
Feature engineering	Preserve clinically meaningful variables while deriving longitudinal and interaction features	Convert raw measurements into risk-relevant predictors such as change in attachment level or cumulative smoking exposure	Supports prediction of both current risk and future worsening
Model selection	Match algorithm complexity to data type, sample size, and interpretability needs	Compare logistic regression, tree-based models, gradient boosting, and neural networks	Identifies the most appropriate model for the clinical prediction task
Model validation	Evaluate discrimination, calibration, external generalizability, and clinical utility	Use internal and multicenter validation before deployment	Prevents premature adoption of poorly generalizable models
Explainability	Provide global and patient-specific explanations	Use interpretable coefficients, feature importance, rule-based logic, or SHAP-type explanations	Helps clinicians understand why a patient or tooth is classified as high risk
Clinical decision support	Embed outputs in dental records or periodontal software	Present risk scores, key drivers, and suggested follow-up intensity during care planning	Converts prediction into actionable periodontal management
Periodic recalibration	Update model performance as populations, workflows, and treatment standards change	Monitor drift and recalibrate with new validated datasets	Maintains accuracy and fairness over time

The final stage is feedback and recalibration after implementation. Risk predictions should be compared with observed progression, tooth loss, treatment response, and maintenance outcomes over time. Long-term studies using staging, grading, and prognostic models show that periodontal prediction must remain dynamic because patient behavior, clinical status, and tooth-level risk can change [19]. The proposed framework therefore treats deployment as the beginning of continuous evaluation rather than the end of model development.

Implementation challenges and future directions

The first implementation challenge is data standardization across clinics, software systems, and imaging protocols. Periodontal probing measurements may vary by examiner, radiographs may differ in angulation and resolution, and behavioral histories may be incompletely recorded. Automated detection of periodontal bone loss can improve scalability, but generalizability across radiographic devices and populations must be demonstrated [9, 21]. Future research should therefore prioritize standardized data dictionaries and multicenter validation cohorts.

The second challenge is missingness and bias in real-world dental data. Patients with irregular attendance may have fewer periodontal charts, fewer radiographs, and less complete behavioral documentation, yet these same patients may be at elevated risk. Predictive models derived from electronic dental records can be powerful, but they may inherit documentation biases from routine practice [33]. The framework therefore requires transparent missing-data strategies and fairness assessment before clinical deployment.

The third challenge is regulatory, ethical, and professional accountability. A periodontal prediction

model may influence recall intervals, treatment intensity, referral decisions, and patient counseling, so errors can have real clinical consequences. Scoping and systematic reviews of artificial intelligence in periodontology stress that implementation must address validation, interpretability, governance, and clinician trust [31, 32]. Dental institutions should therefore develop oversight processes before adopting predictive tools at scale.

Future research should move from retrospective model development toward prospective impact studies. Multicenter cohorts can test whether multimodal prediction improves early intervention, reduces progression, supports maintenance adherence, or prevents tooth loss. Newer data streams, including salivary biomarkers, microbiome profiles, and treatment-response data, may eventually expand the framework beyond clinical, behavioral, and radiographic inputs [13, 30, 35]. The priority, however, should be clinically meaningful validation rather than uncontrolled expansion of data complexity.

Conclusion

This conceptual framework provides an evidence-based blueprint for machine learning-based prediction of periodontal disease progression. It integrates clinical, behavioral, systemic, and radiographic determinants into a structured pathway from data capture to model output and clinical action. Its central contribution is to clarify how heterogeneous periodontal information can be organized for interpretable, clinically useful prognosis.

The framework emphasizes that prediction should support periodontal judgment rather than replace it. Machine learning can help identify high-risk patients and vulnerable teeth, but its outputs must remain transparent, calibrated, and embedded in human decision-making. Clinicians should be able to interpret, question, and contextualize model recommendations within each patient's broader care plan.

Operationalizing this framework will require collaboration among periodontists, radiologists, informaticians, data scientists, software developers, and patients. Future work should validate multimodal models prospectively, evaluate their effect on treatment decisions, and monitor fairness across clinical populations. With careful implementation, machine learning-based periodontal prediction can become a practical tool for personalized prevention, maintenance planning, and long-term tooth preservation.

Acknowledgments: None

Conflict of Interest: None

Financial Support: None

Ethics Statement: None

References

1. Tonetti MS, Greenwell H, Kornman KS. Staging and grading of periodontitis: framework and proposal of a new classification and case definition. J Periodontol. 2018;89(Suppl 1):S159-72. doi:10.1002/JPER.18-0006
2. Papapanou PN, Sanz M, Buduneli N, Dietrich T, Feres M, Fine DH, et al. Periodontitis: consensus report of workgroup 2 of the 2017 World Workshop on the Classification of Periodontal and Peri-Implant Diseases and Conditions. J Periodontol. 2018;89(Suppl 1):S173-82. doi:10.1002/JPER.17-0721
3. Needleman I, Garcia R, Gkranias N, Kirkwood KL, Kocher T, Di Iorio A, et al. Mean annual attachment, bone level, and tooth loss: a systematic review. J Clin Periodontol. 2018;45(Suppl 20):S112-29. doi:10.1111/jcpe.12943
4. Nibali L, Krajewski A, Donos N, Völzke H, Pink C, Kocher T, et al. The effect of furcation involvement on tooth loss in a population without regular periodontal therapy. J Clin Periodontol. 2017;44(8):813-21. doi:10.1111/jcpe.12756
5. Lertpimonchai A, Rattanasiri S, Arj-Ong Vallibhakara S, Attia J, Thakkinstian A. The association between oral hygiene and periodontitis: a systematic review and meta-analysis. Int Dent J. 2017;67(6):332-43. doi:10.1111/idj.12317
6. Leite FRM, Nascimento GG, Scheutz F, López R. Effect of smoking on periodontitis: a systematic review and meta-regression. Am J Prev Med. 2018;54(6):831-41. doi:10.1016/j.amepre.2018.02.014
7. Ravidà A, Troiano G, Qazi M, Saleh MHA, Saleh I, Borgnakke WS, et al. Dose-dependent effect of smoking and smoking cessation on periodontitis-related tooth loss during 10-47 years periodontal maintenance: a retrospective study in compliant cohort. J Clin Periodontol. 2020;47(9):1132-43. doi:10.1111/jcpe.13336
8. Lee JH, Kim DH, Jeong SN, Choi SH. Diagnosis and prediction of periodontally compromised teeth using a deep learning-based convolutional neural network algorithm. J Periodontal Implant Sci.

2018;48(2):114-23. doi:10.5051/jpis.2018.48.2.114

9. Krois J, Ekert T, Meinhold L, Golla T, Kharbot B, Wittemeier A, et al. Deep learning for the radiographic detection of periodontal bone loss. Sci Rep. 2019;9(1):8495. doi:10.1038/s41598-019-44839-3
10. Ravidà A, Qazi M, Troiano G, Saleh MHA, Greenwell H, Kornman K, et al. Using periodontal staging and grading system as a prognostic factor for future tooth loss: a long-term retrospective study. J Periodontol. 2020;91(4):454-61. doi:10.1002/JPER.19-0390
11. Kim J, Lee HS, Song IS, Jung KH. DeNTNet: Deep Neural Transfer Network for the detection of periodontal bone loss using panoramic dental radiographs. Sci Rep. 2019;9(1):17615. doi:10.1038/s41598-019-53758-2
12. Helal O, Göstemeyer G, Krois J, Fawzy El Sayed K, Graetz C, Schwendicke F. Predictors for tooth loss in periodontitis patients: systematic review and meta-analysis. J Clin Periodontol. 2019;46(7):699-712. doi:10.1111/jcpe.13118
13. Huang W, Wu J, Mao YQ, Zhu SW, Huang GF, Petritis B, et al. Developing a periodontal disease antibody array for the prediction of severe periodontal disease using machine learning classifiers. J Periodontol. 2020;91(2):232-43. doi:10.1002/JPER.19-0173
14. Bashir NZ, Rahman Z, Chen SLS. Systematic comparison of machine learning algorithms to develop and validate predictive models for periodontitis. J Clin Periodontol. 2022;49(10):958-69. doi:10.1111/jcpe.13692
15. Chang HJ, Lee SJ, Yong TH, Shin NY, Jang BG, Kim JE, et al. Deep learning hybrid method to automatically diagnose periodontal bone loss and stage periodontitis. Sci Rep. 2020;10(1):7531. doi:10.1038/s41598-020-64509-z
16. Terela H, Strilets V. Supervision and control of labor law compliance in Ukraine under martial law. Ann Organ Cult Leadersh Extern Engagem J. 2024;5:10-6. doi:10.51847/QRKh2QNIig
17. Moore E, Garcia CZ, Ferrari G. A phase II, multicenter, randomized, double-blind study to assess the tolerability of escalating induction doses of Everolimus in patients with metastatic breast cancer (DESIREE). Asian J Curr Res Clin Cancer. 2025;5(1):145-56. doi:10.51847/VReJL5V3r5
18. Ruiz AF, Desta HT, Ismail NS. Oral microbiome dynamics and Surfactant Protein A expression in patients with spontaneous intraoral lesions. J Curr Res Oral Surg. 2025;5(1):176-88. doi:10.51847/81n21UzLVq
19. Tan GY, Lim WM, Ong HX. Perceived control and self-rated health from young adulthood to midlife: findings from the Longitudinal Youth Development Study. Int J Soc Psychol Asp Healthc. 2024;4:100-12. doi:10.51847/bUYUzIFshB
20. Thanathornwong B, Suebnukarn S. Automatic detection of periodontal compromised teeth in digital panoramic radiographs using faster regional convolutional neural networks. Imaging Sci Dent. 2020;50(2):169-74. doi:10.5624/isd.2020.50.2.169
21. Lee CT, Kabir T, Nelson J, Sheng S, Meng HW, Van Dyke TE, et al. Use of the deep learning approach to measure alveolar bone level. J Clin Periodontol. 2022;49(3):260-9. doi:10.1111/jcpe.13574
22. Tsoromokos N, Parinussa S, Claessen F, Moin DA, Loos BG. Estimation of alveolar bone loss in periodontitis using machine learning. Int Dent J. 2022;72(5):621-7. doi:10.1016/j.identj.2022.02.009
23. Chang J, Chang MF, Angelov N, Hsu CY, Meng HW, Sheng S, et al. Application of deep machine learning for the radiographic diagnosis of periodontitis. Clin Oral Investig. 2022;26(11):6629-37. doi:10.1007/s00784-022-04617-4
24. Chen CC, Wu YF, Aung LM, Lin JC, Ngo ST, Su JN, et al. Automatic recognition of teeth and periodontal bone loss measurement in digital radiographs using deep-learning artificial intelligence. J Dent Sci. 2023;18(3):1301-9. doi:10.1016/j.jds.2023.03.020
25. Alqahtani HM, Koroukian SM, Stange K, Schiltz NK, Bissada NF. Identifying factors associated with periodontal disease using machine learning. J Int Soc Prev Community Dent. 2022;12(6):612-22. doi:10.4103/jispcd.JISPCD_188_22
26. Lee CT, Zhang K, Li W, Tang K, Ling Y, Walji MF, et al. Identifying predictors of tooth loss using a rule-based machine learning approach: a retrospective study at university-setting clinics. J Periodontol. 2023;94(10):1231-42. doi:10.1002/JPER.23-0030
27. Guler Ayyildiz B, Karakis R, Terzioglu B, Ozdemir D. Comparison of deep learning methods for the radiographic detection of patients with different periodontitis stages. Dentomaxillofac Radiol. 2024;53(1):32-42. doi:10.1093/dmfr/twad003
28. Rahim-Wöstefeld S, Kronsteiner D, ElSayed S, ElSayed N, Eickholz P, Pretzl B. Development of a prognostic tool: based on risk factors for tooth

loss after active periodontal therapy. Clin Oral Investig. 2022;26(1):813-22. doi:10.1007/s00784-021-04060-x

29. Feher B, de Souza Oliveira EH, Mendes Duarte P, Werdich AA, Giannobile WV, Feres M. Machine learning-assisted prediction of clinical responses to periodontal treatment. J Periodontol. 2025;96(11):1199-212. doi:10.1002/JPER.24-0737
30. Polizzi A, Quinzi V, Lo Giudice A, Marzo G, Leonardi R, Isola G. Accuracy of artificial intelligence models in the prediction of periodontitis: a systematic review. JDR Clin Trans Res. 2024;9(4):312-24. doi:10.1177/23800844241232318
31. Scott J, Biancardi AM, Jones O, Andrew D. Artificial intelligence in periodontology: a scoping review. Dent J (Basel). 2023;11(2):43. doi:10.3390/dj11020043
32. Swinckels L, de Keijzer A, Loos BG, Applegate RJ, Kookal KK, Kalenderian E, et al. A personalized periodontitis risk based on nonimage electronic dental records by machine learning. J Dent. 2025;153:105469. doi:10.1016/j.jdent.2024.105469
33. Lee CT, Zhang K, Li W, Tang K, Ling Y, Walji MF, et al. Identifying predictors of the tooth loss phenotype in a large periodontitis patient cohort using a machine learning approach. J Dent. 2024;144:104921. doi:10.1016/j.jdent.2024.104921
34. Deng K, Zonta F, Yang H, Pelekos G, Tonetti MS. Development of a machine learning multiclass screening tool for periodontal health status based on non-clinical parameters and salivary biomarkers. J Clin Periodontol. 2024;51(12):1547-60. doi:10.1111/jcpe.13856
35. Enevold C, Nielsen CH, Christensen LB, Kongstad J, Fiehn NE, Hansen PR, et al. Suitability of machine learning models for prediction of clinically defined Stage III/IV periodontitis from questionnaires and demographic data in Danish cohorts. J Clin Periodontol. 2024;51(12):1561-73. doi:10.1111/jcpe.13874
36. Santamaria P, Troiano G, Serroni M, Araùjo TG, Ravidà A, Nibali L. Exploring the accuracy of tooth loss prediction between a clinical periodontal prognostic system and a machine learning prognostic model. J Clin Periodontol. 2024;51(10):1333-41. doi:10.1111/jcpe.14023
37. Wong Y, Lin S, Cheng H, Hsieh T, Hsiue T, Chung H, et al. Understanding the impact of medical humanities on internship training and performance. Ann Pharm Educ Saf Public Health Advocacy. 2025;5:12-21. doi:10.51847/Z1fogzPksy
38. Min-Ho L, Jin-Woo C. Genomic landscape and clinical implications of MTAP loss in advanced gastrointestinal malignancies. Arch Int J Cancer Allied Sci. 2024;4(1):67-80. doi:10.51847/1wgE2vWBFG
39. Mitchell F, Erio K, Sohal S. Ethical considerations in adolescent participation in vaccine trials: qualitative perspectives on family decision-making from a human papillomavirus vaccine trial in Tanzania. Asian J Ethics Health Med. 2024;4:107-24. doi:10.51847/SURdqWS9om